



Publication Year	2020
Acceptance in OA	2021-11-16T15:47:19Z
Title	The accretion history of high-mass stars: An ArTéMiS pilot study of Infrared Dark Clouds
Authors	Peretto, N., Rigby, A., André, Ph., Könyves, V., Fuller, G., Zavagno, A., Schuller, F., Arzoumanian, D., Bontemps, S., Csengeri, T., Didelon, P., Duarte-Cabral, A., Palmeirim, P., PEZZUTO, Stefano, Revéret, V., Roussel, H., Shimajiri, Y.
Publisher's version (DOI)	10.1093/mnras/staa1656
Handle	http://hdl.handle.net/20.500.12386/31092
Journal	MONTHLY NOTICES OF THE ROYAL ASTRONOMICAL SOCIETY
Volume	496

The accretion history of high-mass stars: an ArTéMiS pilot study of infrared dark clouds

N. Peretto¹,¹★ A. Rigby,¹ Ph. André,² V. Könyves,³ G. Fuller,⁴ A. Zavagno,^{5,6}
F. Schuller^{2,7}, D. Arzoumanian,⁸ S. Bontemps,⁹ T. Csengeri,⁹ P. Didelon,²
A. Duarte-Cabral¹, P. Palmeirim⁸, S. Pezzuto¹⁰, V. Revéret,² H. Roussel¹¹
and Y. Shimajiri¹²

¹Cardiff University, School of Physics & Astronomy, Queen's buildings, The parade, Cardiff CF24 3AA, UK

²Laboratoire d'Astrophysique (AIM), CEA/DRF, CNRS, Université Paris-Saclay, Université Paris Diderot, Sorbonne Paris Cité, F-91191 Gif-sur-Yvette, France

³Jeremiah Horrocks Institute, University of Central Lancashire, Preston PR1 2HE, UK

⁴Jodrell Bank Centre for Astrophysics, Department of Physics and Astronomy, The University of Manchester, Oxford Road, Manchester M13 9PL, UK

⁵Aix Marseille Univ, CNRS, CNES, LAM, Marseille, France

⁶Institut Universitaire de France

⁷Leibniz-Institut für Astrophysik Potsdam (AIP), An der Sternwarte 16, D-14482 Potsdam, Germany

⁸Instituto de Astrofísica e Ciências do Espaço, Universidade do Porto, CAUP, Rua das Estrelas, P-4150-762 Porto, Portugal

⁹Laboratoire d'Astrophysique de Bordeaux, Univ. Bordeaux, CNRS, B18N, allée Georoy Saint-Hilaire, F-33615 Pessac, France

¹⁰INAF, IAPS Via fosse del Cavaliere 100, I-00133 Roma, Italy

¹¹Institut d'Astrophysique de Paris, Sorbonne Université, CNRS (UMR7095), F-75014 Paris, France

¹²National Astronomical Observatory of Japan, Osawa 2-21-1, Mitaka, Tokyo 181-8588, Japan

Accepted 2020 June 7. Received 2020 June 5; in original form 2020 April 14

ABSTRACT

The mass growth of protostars is a central element to the determination of fundamental stellar population properties such as the initial mass function. Constraining the accretion history of individual protostars is therefore an important aspect of star formation research. The goal of the study presented here is to determine whether high-mass (proto)stars gain their mass from a compact (<0.1 pc) fixed-mass reservoir of gas, often referred to as dense cores, in which they are embedded, or whether the mass growth of high-mass stars is governed by the dynamical evolution of the parsec-scale clump that typically surrounds them. To achieve this goal, we performed a $350\text{-}\mu\text{m}$ continuum mapping of 11 infrared dark clouds, along side some of their neighbouring clumps, with the ArTéMiS camera on APEX. By identifying about 200 compact ArTéMiS sources, and matching them with *Herschel* Hi-GAL $70\text{-}\mu\text{m}$ sources, we have been able to produce mass versus temperature diagrams. We compare the nature (i.e. starless or protostellar) and location of the ArTéMiS sources in these diagrams with modelled evolutionary tracks of both core-fed and clump-fed accretion scenarios. We argue that the latter provide a better agreement with the observed distribution of high-mass star-forming cores. However, a robust and definitive conclusion on the question of the accretion history of high-mass stars requires larger number statistics.

Key words: accretion – stars: protostars – stars: formation.

1 INTRODUCTION

Our knowledge about star formation has made tremendous progress in the past few years (André et al. 2014; Motte, Bontemps & Louvet 2018b). As a result of the science exploitation of *Herschel* data,

our picture of how matter is condensed from diffuse clouds to stars has been significantly improved. In particular, one striking result has been the ubiquity of interstellar filaments in all types of molecular clouds, including low-mass star-forming (André et al. 2010), massive-star-forming (Molinari et al. 2010), and even non-star-forming clouds (Men'shchikov et al. 2010). In nearby, and mostly low-mass, star-forming regions, nearly all the dense gas ($n_{\text{H}_2} \geq 10^4 \text{ cm}^{-3}$; $N_{\text{H}_2} \geq 10^{22} \text{ cm}^{-2}$) is concentrated within 0.1-

* E-mail: nicolas.peretto@astro.cf.ac.uk

pc-width filaments (Arzoumanian et al. 2011, 2019). Moreover, the vast majority (~ 75 percent) of pre-stellar cores are found to lie within these filaments (Könyves et al. 2015, 2020; Ladjelate et al. 2020). It has therefore been proposed that solar-type star-forming cores form as the result of Jeans-type gravitational instabilities developing along self-gravitating filaments (Inutsuka & Miyama 1997), providing a compact and, at least to first order, fixed mass reservoir for the protostars forming inside them. As a result, such protostars are said to be *core-fed*. This scenario is believed to explain the shape of the pre-stellar core mass function as observed in a number of low-mass star-forming clouds (e.g. Motte, Andre & Neri 1998; Könyves et al. 2015) and therefore the origin of the base of the initial mass function (IMF) from ~ 0.1 to $5 M_{\odot}$ (cf. André et al. 2014, 2019; Lee, Hennebelle & Chabrier 2017).

At the high-mass end of the IMF, however, it is well known that thermal Jeans-type fragmentation cannot explain the formation of cores more massive than a few solar masses (e.g. Bontemps et al. 2010). In fact, the search for compact high-mass pre-stellar cores with ALMA systematically reveals that reasonable candidates identified with single-dish telescopes are systematically sub-fragmented into low-mass cores (e.g. Louvet et al. 2019; Sanhueza et al. 2019; Svoboda et al. 2019). Therefore, the formation of massive stars requires additional physics. One key difference between low- and high-mass star formation is that self-gravitating mass reservoirs in massive star-forming regions are larger (in mass and size) by several orders of magnitude (e.g. Beuther et al. 2013). These parsec-scale structures are often referred to as *clumps*. Even though the question about how massive stars form is still very much debated, observations (e.g. Peretto, André & Belloche 2006; Schneider et al. 2010; Duarte-Cabral et al. 2013; Peretto et al. 2013; Urquhart et al. 2014; Csengeri et al. 2017) and simulations (e.g. Bonnell, Vine & Bate 2004; Smith, Longmore & Bonnell 2009; Wang et al. 2010; Vázquez-Semadeni et al. 2019) converge towards a picture where massive stars form at the centre of globally collapsing clumps, quickly growing in mass as a result of large infall rates ($\dot{m}_{\text{inf}} \sim 10^{-3} M_{\odot} \text{ yr}^{-1}$). In this picture, massive stars are said to be *clump-fed*. Two questions naturally arise: (i) Is the clump-fed scenario the dominant mode of high-mass star formation? (ii) If yes, then around what core mass does the transition from core-fed to clump-fed star formation scenarios occur?

Constraining the process through which stars gain mass is a major goal of star formation research. A number of studies have provided predictions regarding the mass and luminosity functions of protostars in the context of both the *core-fed* and *clump-fed* scenarios (Myers 2009, 2012; McKee & Offner 2010; Offner & McKee 2011). Such models have been compared to observations of nearby, *low-mass*, protoclusters, but flux uncertainties, the limited protostellar mass range, and the low-number statistics have so far prevented model discrimination (with the exception of the Single Isothermal Sphere model, which is inconsistent with observations – Offner & McKee 2011). Mass-luminosity diagrams of protostellar cores have also been often used to constrain the time evolution of protostars. Theoretical evolutionary tracks have been computed, mostly assuming a fixed initial mass reservoir, i.e. *core-fed* (Bontemps et al. 1996; Andre, Ward-Thompson & Barsony 2000; André et al. 2008; Molinari et al. 2008; Duarte-Cabral et al. 2013).

Recently, the most complete sample of clumps in the Galactic Plane has been identified using the *Herschel* Hi-GAL survey (Elia et al. 2017) and such tracks were used to constrain the time evolution

of the Hi-GAL parsec-scale clumps. Even though the number statistics are impressive (with more than 10^5 sources), the lack of angular resolution prevents us from probing the evolution of individual dense cores, the progenitors of single/small systems of stars. The ALMAGAL project (PIs: A. MOLINARI, P. SCHILKE, C. BATTERSBY, P. HO), i.e. the follow-up at high-angular resolution of ~ 1000 Hi-GAL sources with ALMA, will provide in the near future the first large sample of individual protostellar cores. But even then, the selection bias towards the massive-star-forming clumps will likely prevent answering the question about the transition regime between high-mass and low-mass accretion scenarios. Duarte-Cabral et al. (2013) have constrained the time evolution of a sample of massive-star-forming cores observed at high angular resolution with the IRAM PdBI. In that study, the authors compute the M_{core} versus L_{bol} and F_{CO} versus L_{bol} evolution using simple core evolution models and use Monte Carlo simulations to populate the diagram. While very promising, the low-number statistics of nine cores, and the focus on massive protostellar sources, limit the possibility of constraining core mass growth scenarios across all evolutionary stages and masses.

In this paper, we present new ArTéMiS/APEX 350- μm continuum observations of a sample of infrared dark clouds (IRDCs hereafter). The observed fields are large (the largest presented here is $\sim 400 \text{ arcmin}^2$ – see Fig. 1) allowing us to get a complete census of the source population within the targeted regions at relatively high angular resolution (i.e. $\sim 8''$). This is the complementary approach to surveys such as Hi-GAL and ALMAGAL. The main goal of this study is to demonstrate the potential of ArTéMiS in determining the relative importance of *core-fed* and *clump-fed* scenarios in the context of high-mass star formation. This work will serve as a pilot study for CAFFEINE, an ArTéMiS large programme currently underway.

2 TARGETS AND OBSERVATIONS

We targeted a total of 11 IRDCs from the Peretto & Fuller (2009) catalogue, selected to have a H_2 column density peak above 10^{23} cm^{-2} and located at very similar distances, i.e. $2.6 \pm 0.3 \text{ kpc}$, with the exception of SDC326.796 + 0.386 (see Table 1). Kinematic distances to all sources, but one, have been estimated using the MALT90 $\text{N}_2\text{H}^+(1-0)$ data (Foster et al. 2013) and the Reid et al. (2009) and Reid et al. (2014) galactic rotation model. Because SDC340.928-1.042 was not mapped by MALT90, we used the ThrUMMS $^{13}\text{CO}(1-0)$ data instead (Barnes et al. 2015) to obtain its systemic velocity. These data show that it is part of the same molecular cloud as SDC340.969-1.020 and we therefore assigned the same velocity to both IRDCs. For all sources, we adopted the near heliocentric distance as most IRDCs are located at the near distance (Ellsworth-Bowers et al. 2013; Giannetti et al. 2015). When more than one ArTéMiS clumps are part of a single IRDC, we checked that the velocity and corresponding distances of individual clumps are similar, which turned to be always the case. The typical uncertainty on these distances is 15 percent (Reid et al. 2009). Table 1 shows the main properties of the sources, including their effective radii and background-subtracted masses as estimated from the *Herschel* column density maps from Peretto et al. (2016) within a H_2 column density contour level of $2 \times 10^{22} \text{ cm}^{-2}$. The 11 IRDCs have been mapped as part of five individual fields that in some cases (in particular for the largest of all, i.e. the SDC326 field) include extra sources. For all of these extra sources, we ensured that their kinematic distances were similar to the average field distance by using the same method as described above.




Figure 1. 350- μm ArTéMiS image of the SDC326 field (in logarithmic scale), that includes four targeted IRDCs whose ID numbers (in cyan) correspond to those indicated in Table 1. The area within the red polygon roughly corresponds to the area of uniform noise. The 10-pc bar scale at the bottom of the image assumes a distance of 2.6 kpc.

Table 1. Target properties. The V_{lsr} column corresponds to velocities that have been obtained using the MALT90 $\text{N}_2\text{H}^+(1-0)$ line, with the exception of SDC340.928-1.042 (marked with an asterisk) for which we used the ThrUMMS $^{13}\text{CO}(1-0)$ line (see text). IRDCs having several velocities/distances correspond to ArTéMiS maps where several clumps have been detected. Noise values correspond to the median rms noise in each of the five observed fields, for both the original ArTéMiS images and the Gaussian filtered ones.

ID#	IRDC name	Coordinates (J2000)	V_{lsr} (km/s)	Distance (kpc)	Radius (pc)	Mass (M_{\odot})	Unfilt. noise (Jy beam^{-1})	Filt. noise (Jy beam^{-1})
1	SDC326.476 + 0.706	15:43:16.4 – 54:07:13	–40.5	2.7	1.08	3730	0.48	0.26
2	SDC326.611 + 0.811	15:43:36.3 – 53:57:45	–37.0	2.5	1.29	3260	0.48	0.26
3	SDC326.672 + 0.585	15:44:57.3 – 54:07:14	–41.3	2.7	0.91	4120	0.48	0.26
4	SDC326.796 + 0.386	15:46:20.9 – 54:10:44	–20.4	1.6	0.42	240	0.48	0.26
5	SDC328.199 – 0.588	15:57:59.6 – 53:58:01	–44.3 – 38.7	2.9 2.6	2.77	33220	0.63	0.41
6	SDC340.928 – 1.042	16:55:01.4 – 45:11:42	–24.1*	2.3	0.73	640	0.46	0.31
7	SDC340.969 – 1.020	16:54:57.1 – 45:09:04	–24.1	2.3	0.66	2630	0.46	0.31
8	SDC343.722 – 0.178	17:00:49.6 – 42:26:05	–28.0 – 26.7 – 25.6	2.8 2.7 2.6	1.42	5270	0.51	0.38
9	SDC343.735 – 0.110	17:00:32.6 – 42:25:02	–27.3	2.7	0.45	510	0.51	0.38
10	SDC343.781 – 0.236	17:01:13.0 – 42:27:42	–27.1	2.5	0.46	360	0.51	0.38
11	SDC345.000 – 0.232	17:05:10.8 – 41:29:08	–27.8	2.9	2.14	16160	0.43	0.30

All targets were observed at 350 μm with APEX and the ArTéMiS camera¹ (Revéret et al. 2014; André et al. 2016) between September 2013 and August 2014 (Onsala projects O-091.F-9301A and O-

093.F-9317A). The angular resolution at 350 μm with APEX is $\theta_{\text{beam}} = 8''$. Observations have been carried out with individual maps of $6'' \times 6''$, with a minimum of two coverages per field with different scanning angles. The scanning speed ranged from 20''/s to 3''/s and the cross-scan step between consecutive scans from 6'' to 12''. The 350- μm sky opacity (at zenith) and precipitable water vapour at the telescope were typically between 0.7 and 1.9 and

¹Note that at the time of these observations, the 450- μm array was not available.

between 0.35 mm and 0.85 mm, respectively. Absolute calibration was achieved by observing Mars as a primary calibrator, with a corresponding calibration uncertainty of ~ 30 per cent. Regular calibration and pointing checks were performed by taking short spiral scans towards the nearby secondary calibrators B13134, IRAS 16293, and G5.89 every ~ 0.5 -1.0 h. The pointing accuracy was within $\sim 3''$. Data reduction was performed using the APIS pipeline running in IDL.² The ArTéMiS images can be seen in Fig. 1 and in Appendix A.

3 COMPACT SOURCE IDENTIFICATION

In order to identify compact sources in all our fields, we first convolved all ArTéMiS images with a Gaussian of FWHM of 0.2 pc ($\sim 15''$ at 2.6 kpc) and subtracted that convolved image from the original image. By doing so, we filter our ArTéMiS images from emission on spatial scales ≥ 0.2 pc, and the comparison between sources becomes independent of their background properties. We then identify compact sources using dendrograms (e.g. Rosolowsky et al. 2008; Peretto & Fuller 2009) on signal-to-noise ratio (SNR) maps (see Fig. 2). For that purpose, we computed noise maps, σ_{map} , from the ArTéMiS weight maps, ω_{map} (proportional to the integration time at every position in the map), and a noise calibration, σ_{ref} , estimated on an emission-free area of the filtered ArTéMiS maps following: $\sigma_{\text{map}} = \sigma_{\text{ref}} \sqrt{(\omega_{\text{ref}}/\omega_{\text{map}})}$, where ω_{ref} is the average weight estimated in the same region as σ_{ref} . The calibration σ_{ref} is computed on the Gaussian filtered images (see Table 1 for median rms noise values). Our dendrogram source identification uses a starting level of $2\sigma_{\text{map}}$, a step of $3\sigma_{\text{map}}$ (i.e. all sources must have a minimum SNR peak of 5), and a minimum source solid angle of 50 per cent of the beam solid angle that translates into a minimum effective diameter of $\sim 5.6''$ (~ 0.07 pc at a distance of 2.6 kpc), i.e. 70 per cent of the beam FWHM. The *leaves* of the dendrogram (i.e. structures that exhibit no further fragmentation within the boundaries set by the input parameters of the extraction) are then used as masks in the filtered ArTéMiS images to measure the peak flux density of every source. In the context of the present study, this is the only parameter we are interested in (see Section 5.2). As it can be seen in, e.g. Fig. 1, the noise in the image is non-uniform and increases towards the edge of the image. In order to reduce the potential bias in the source detection created by a non-uniform noise, we defined, by hand and for each field, a mask that cuts out the noisy edges. In the following, we consider only the sources that fall within this mask. In total, across all fields, we detect 203 compact ArTéMiS sources. Table 2 provides information on individual sources, and individual cutout images of each source can be found in Appendix C. Note that the source extraction parameters used in this paper are rather conservative and as a result faint sources might remain unidentified. However, the non-detection of such sources does not affect any of the results discussed here.

4 ASSOCIATIONS OF ARTÉMiS SOURCES WITH HI-GAL AND ATLASGAL SOURCE CATALOGUES

In the past 10 yr, far-IR and (sub-)millimetre continuum surveys of the Galactic plane have significantly contributed to improve

our knowledge of massive star formation (Schuller et al. 2009; Molinari et al. 2010; Aguirre et al. 2011; Moore et al. 2015). However, even though these surveys have been, and still are, rich sources of information regarding massive star formation studies, one key issue is the lack of high-resolution, high-sensitivity observations of the cold dust on similar angular resolution as the Herschel 70- μm band ($\sim 7''$ resolution), which traces the protostars' luminosities. By filling in this gap, ArTéMiS observations allow us to unambiguously determine the envelope mass of young protostellar objects throughout the Galactic plane. In order to demonstrate the advancement that sensitive ArTéMiS observations provide over existing surveys, we here compare sub-millimetre source detections with Hi-GAL (Elia et al. 2017), and ATLASGAL (Csengeri et al. 2014), along with performing a *Herschel* 70- μm source association using the Molinari et al. (2016) catalogue.

Association between our ArTéMiS sources and sources in published catalogues is performed by searching sources whose published coordinates lie within one beam of the central coordinates of the ArTéMiS source. We therefore used an angular separation of $8''$ when performing the 70- μm association, $19''$ when performing the association with ATLASGAL sources, and $36''$ when performing the Hi-GAL clump association. The statistics of the number of sources within each field and their respective association with ArTéMiS sources are given in Tables 3 and 4. These statistics show a number of important points. First, 14 per cent of the ArTéMiS sources are newly identified sources that do not belong to any of the three catalogues we searched for. Also, about 54 per cent of Hi-GAL clumps and 63 per cent of ATLASGAL sources have an ArTéMiS detection associated with them. Finally, about 42 per cent of the ArTéMiS sources have a published 70- μm source associated with them, but when looking at the individual cutouts provided in Appendix C, one realizes that an extra ~ 25 per cent of sources have locally peaked 70- μm or 8- μm emission towards them. This means that about 67 per cent of the ArTéMiS sources are protostellar, and about 33 per cent are starless (down to the 70- μm sensitivity of Hi-GAL of ~ 0.1 Jy – Molinari et al. 2016).

Fig. 3 shows examples for each association type (see also Appendices B and C). In this figure, we display seven sources, four of which are detected with ArTéMiS and three of which are not. We also show these seven sources at different wavelengths in order to better understand the type of sources that we do, and do not, detect with ArTéMiS. On the same figure, the symbols indicate when a source has been identified in the three different source catalogues used. By looking at Fig. 3, it becomes clear that ArTéMiS is particularly good at identifying protostellar sources. In fact, even the source in the fourth column, which has not been identified in any of the three catalogue used, and which has therefore no *Herschel* 70- μm entries in the Molinari et al. (2016) catalogue, seems to be associated with a faint point-like 70- μm emission (as mentioned above, ~ 25 per cent of sources fall in this category of sources). On the other hand, all three sources displayed in Fig. 3 that have not been detected with ArTéMiS have no 70- μm emission associated with them. The source in the fifth column is clearly seen in the ArTéMiS data but falls just below our 5σ threshold of detection. In a similar way as displayed in Fig. 3, we looked at all individual ArTéMiS sources we identified to ensure the quality of the detection. Individual images of each ArTéMiS source can be found in Appendix C.

²http://www.apex-telescope.org/instruments/pi/artemis/data_reduction/




Figure 2. Left: 350- μm ArTéMiS image of part of SDC326.611 + 0.811 field (linear scale). Contours start at 1 Jy beam^{-1} by steps of 1 Jy beam^{-1} . The green circles show the positions of *Herschel* clumps from Elia et al. (2017), while the magenta squares show the positions of ATLASGAL clumps from Csengeri et al. (2014). Middle: Gaussian filtered image at 0.2 pc of the same source. Contours start at 0.5 Jy beam^{-1} in steps of 0.5 Jy beam^{-1} . Right: Signal-to-noise ratio map of the same source. Contours start at 2 by steps of 3. The yellow crosses show the position of the ArTéMiS sources, while the cyan circles show the positions of 70- μm sources from Molinari et al. (2016).

Table 2. Properties of the first 10 ArTéMiS sources identified in the SDC326 field. First column: Source ID number; second column: Galactic longitude; third column: Galactic latitude; fourth column: Original ArTéMiS peak flux density (i.e. non-filtered) and associated uncertainties; fifth column: Filtered ArTéMiS peak flux density and associated uncertainties; sixth column: Dust temperature estimated on 0.1-pc scale (see Section 6.1) and associated uncertainties; seventh column: Gas mass estimated on 0.1-pc scale (see Section 6.1) and associated uncertainties; eighth column: Internal luminosity and associated uncertainties. If a value is given, it means that the ArTéMiS source has a *Herschel* 70- μm source from Molinari et al. (2016) associated with it; ninth column: Is there a *Herschel* clump from Elia et al. (2017) associated with it? ‘y’ for yes, ‘n’ for no; tenth column: Is there an ATLASGAL source from Csengeri et al. (2014) associated with it? ‘y’ for yes, ‘n’ for no; and eleventh column: Can we visually identify a mid-infrared (70- μm and/or 8- μm peak) peak on the individual cutout images in Appendix C? ‘y’ for yes, ‘n’ for no. The full table can be found online.

ID #	l (degree)	b (degree)	S_{ν}^{pk} (Jy beam^{-1})	$S_{\nu}^{\text{pk}}[\text{filt}]$ (Jy beam^{-1})	$T_{\text{dust}}(0.1\text{pc})$ (K)	$M_{\text{gas}}(0.1\text{pc})$ (M_{\odot})	L_{int} ($\times 10^3 L_{\odot}$)	H clump?	A clump?	mid-IR?
SDC326										
Field										
1	326.7951	0.3817	17.4 ± 5.2	9.8 ± 2.9	35.1 ± 7.0	$3.8^{+2.9}_{-1.6}$	$1.05^{+0.26}_{-0.32}$	y	y	y
2	326.6328	0.5204	3.9 ± 1.2	1.4 ± 0.4	30.3 ± 6.1	$2.0^{+1.6}_{-0.9}$	$0.33^{+0.10}_{-0.10}$	y	n	y
3	326.6336	0.5288	2.6 ± 0.8	1.0 ± 0.3	30.4 ± 6.1	$1.4^{+1.1}_{-0.6}$	–	y	y	n
4	326.6577	0.5104	6.7 ± 2.0	2.6 ± 0.8	30.7 ± 6.1	$3.6^{+2.8}_{-1.6}$	$0.63^{+0.19}_{-0.19}$	y	n	y
5	326.6622	0.5200	24.5 ± 7.4	11.6 ± 3.5	37.1 ± 7.4	$11.0^{+8.3}_{-4.6}$	$9.41^{+2.64}_{-2.80}$	y	y	y
6	326.6584	0.5169	13.8 ± 4.1	3.4 ± 1.0	25.1 ± 5.0	$6.5^{+5.4}_{-3.0}$	–	y	y	n
7	326.6345	0.5328	3.3 ± 1.0	1.6 ± 0.5	34.6 ± 6.9	$1.7^{+1.4}_{-0.7}$	$0.87^{+0.24}_{-0.26}$	y	y	y
8	326.5636	0.5873	2.5 ± 0.8	0.9 ± 0.3	22.4 ± 4.5	$2.5^{+2.1}_{-1.2}$	–	n	n	y
9	326.6857	0.4950	3.9 ± 1.2	1.9 ± 0.6	28.2 ± 5.6	$3.0^{+2.8}_{-1.4}$	$0.21^{+0.07}_{-0.06}$	y	y	y
10	326.6272	0.5525	1.9 ± 0.6	1.2 ± 0.4	35.0 ± 7.0	$1.3^{+1.1}_{-0.6}$	$0.77^{+0.25}_{-0.23}$	y	n	y

5 PHYSICAL PROPERTIES OF ARTÉMIS SOURCES

5.1 Dust temperatures

A key characteristic of the compact sources we identified within our ArTéMiS data is their dust temperature. Dust temperatures are needed to estimate the mass of these sources but also can be used as an evolutionary tracer of the sources as dust tends to become warmer as star formation proceeds. Here we have computed dust temperatures in two different ways.

5.1.1 Far-infrared colour temperature, T_{col}

In order to compute dust temperatures of interstellar structures, one usually needs multiwavelength observations to get a reasonable coverage of the spectral energy distribution. One problem we are facing is the lack of complementary far-IR sub-millimetre observations at similar angular resolution to our ArTéMiS data. *Herschel* observations represent the best data set available regarding the characterization of cold interstellar dust emission. However, at 250 μm , the angular resolution of *Herschel* is ~ 2.5 times worse than that of APEX at 350 μm . Another big difference between the

Table 3. ArTéMiS source association statistics with Hi-GAL and ATLASGAL sources from Molinari et al. (2016), Elia et al. (2017), and Csengeri et al. (2014) catalogues. Provides the number of ArTéMiS sources in each field, and how many are associated with at least one Hi-GAL clump, one Hi-GAL 70- μ m source, and one ATLASGAL source (see text). The last column provides the number of sources with no association from any of these three catalogues. The bottom line gives the summary across all fields.

Fields	# ArTéMiS sources	# ArTéMiS with Hi-GAL clumps	# ArTéMiS with Hi-GAL 70 μ m	# ArTéMiS with ATLASGAL	# ArTéMiS no association
SDC326	129	104	52	42	19
SDC328	31	24	13	9	5
SDC340	11	8	4	6	2
SDC343	13	12	11	9	1
SDC345	19	18	6	8	1
ALL	203	166	86	74	28

Table 4. Hi-GAL and ATLASGAL source association statistics [taken from Molinari et al. (2016), Elia et al. (2017), and Csengeri et al. (2014) catalogues] with ArTéMiS sources. Provides the number of Hi-GAL clumps, Hi-GAL 70- μ m sources, and ATLASGAL sources in each field, and how many are associated with at least one ArTéMiS sources. The bottom line gives the summary across all fields.

Fields	# Hi-GAL clump in field	# Hi-GAL clumps with ArTéMiS	# Hi-GAL 70 μ m in field	# Hi-GAL 70 μ m with ArTéMiS	# ATLASGAL in field	# ATLASGAL with ArTéMiS
SDC326	87	43	275	52	37	22
SDC328	20	13	114	13	14	8
SDC340	12	5	38	4	8	5
SDC343	12	8	35	11	9	8
SDC345	19	12	43	6	10	6
ALL	150	81	505	86	78	49




Figure 3. Examples of association types. Each column represents a different source, the first four being ArTéMiS detections (from left to right: SDC326 #123, #126, #127, and #118), the last three being *Herschel* and/or ATLASGAL detections. Each row represents a given wavelength, from top to bottom: ArTéMiS 350 μ m; filtered ArTéMiS 350 μ m; ATLASGAL 870 μ m; *Herschel* 250 μ m; *Herschel* 160 μ m; *Herschel* 70 μ m; and *Spitzer* 8 μ m. The different symbols indicate if the source appears in a given catalogue: Crosses for ArTéMiS detections; large circles for *Herschel* clumps; small circles for *Herschel* 70 μ m detections; and squares for ATLASGAL detections.

two data sets is that *Herschel* is sensitive to all spatial scales and therefore recovers a lot more diffuse structures than within our ArTéMiS data. Here, we use the ratio between the 160- μ m and 250- μ m *Herschel* intensities at the location of each ArTéMiS source as a

measure of the source dust temperature (Peretto et al. 2016). In that respect, we first need to measure the local background intensities of each source. We do this by measuring the minimum 250- μ m intensity value within an annulus surrounding each of the ArTéMiS source, along with the corresponding 160- μ m intensity at the same position. The reason behind choosing the lowest 250- μ m intensity is that the local background around these sources can be complex and made of other compact sources, filaments, etc. Therefore, taking, as it is often done, an average of the intensities within the annulus would result in an uncertain background intensity estimate. By focusing on the single faintest 250- μ m pixel, we are relatively confident to take the background at the lowest column density point within the annulus, which should provide a reasonable estimate of the local background of the compact sources we are interested in. We finally subtract the local background measurements from the measured 250- μ m and 160- μ m peak intensities within the source mask. The resulting background-subtracted fluxes are used to compute the far-infrared colour dust temperatures of each ArTéMiS sources (Peretto et al. 2016).

5.1.2 Internal temperature, T_{int}

For a spherical protostellar core, in the situation where dust emission is optically thin, and where the bulk of the source luminosity is in the far-infrared, one can show that flux conservation leads to the following temperature profile (Terebey, Chandler & Andre 1993):

$$T_{\text{int}} = T_0 \left(\frac{r}{r_0} \right)^{-2/(\beta+4)} \left(\frac{L_{\text{int}}}{L_0} \right)^{1/(\beta+4)}, \quad (1)$$

where β is the spectral index of the specific dust opacity law, and (T_0 , r_0 , L_0) are normalization constants and, following Terebey et al. (1993), are here set to (25 K, 0.032 pc, 520 L_{\odot}), respectively. By integrating over the volume of the core, and assuming a given volume density profile, one can then obtain an expression for the

mass-averaged temperature \bar{T}_{int} . Here, we assume that $\rho \propto r^{-2}$, which leads to the following relation:

$$\bar{T}_{\text{int}} = \left(\frac{\beta + 4}{\beta + 2} \right) T_{\text{int}}. \quad (2)$$

Given the luminosity of the source, one can then compute the average dust temperature within a given radius r . In order to compute the bolometric luminosities of ArTéMiS sources, we exploit their tight relationship with 70- μm fluxes (Dunham et al. 2008; Ragan et al. 2012; Elia et al. 2017). Here, each ArTéMiS source has been checked against the Molinari et al. (2016) 70- μm source catalogue (see Section 4) and their corresponding 70- μm fluxes come from the same catalogue. Then, we convert fluxes into luminosities using the following relation (Elia et al. 2017):

$$L_{\text{int}} = 25.6 \left(\frac{F_{70\ \mu\text{m}}}{10\text{Jy}} \right) \left(\frac{d}{1\text{kpc}} \right)^2 L_{\odot}, \quad (3)$$

where $F_{70\ \mu\text{m}}$ is the 70- μm flux of the ArTéMiS source in Jy. This relation is very similar to that obtained for low-mass protostellar objects by Dunham et al. (2008):

$$L_{\text{int}} = 20.0 \left(\frac{F_{70\ \mu\text{m}}}{10\text{Jy}} \right)^{0.94} \left(\frac{d}{1\text{kpc}} \right)^{1.88} L_{\odot}. \quad (4)$$

Since we are using the same *Herschel* data sets as in Elia et al. (2017), we here use the former relationship. Note that these authors have identified a third relation between L_{int} and $F_{70\ \mu\text{m}}$ for sources that do not have a known *Spitzer* or *WISE* mid-infrared at 24 μm or 21 μm , respectively. However, for simplicity, we here use only equation (3), the dependence of T_{int} on L_{int} is in any case very shallow. Finally, by plugging in the corresponding luminosities in equation (1), and by setting $\beta = 2$ (e.g. Hildebrand 1983), we can obtain \bar{T}_{int} for every ArTéMiS protostellar source.

5.1.3 Comparison between T_{col} and \bar{T}_{int}

Our estimates of T_{col} and \bar{T}_{int} use independent *Herschel* data and make use of different sets of assumptions to compute the same quantity, i.e. the dust temperature of ArTéMiS sources. In order to decide which of these two sets of temperatures is the most appropriate to use, we plotted them against each other (see Fig. 4). This can be done only for ArTéMiS sources with an associated 70- μm source. For the purpose of making Fig. 4, \bar{T}_{int} has been here estimated within a radius equivalent to the *Herschel* 250- μm beam (i.e. 0.23 pc at 2.6-kpc distance) so that the comparison remains valid. Uncertainties have been estimated by using Monte Carlo propagation. Uncertainties for \bar{T}_{int} are much lower as a result of its shallow dependence on $F_{70\ \mu\text{m}}$. One can see that the two sets of values are well correlated to each other, with a median ratio $\bar{T}_{\text{int}}/T_{\text{col}} = 1.28^{+0.20}_{-0.25}$. This shows that, for most of the points in Fig. 4, the far-IR colour temperature is lower by ~ 28 per cent compared to its internal temperature counterpart. Interestingly, Peretto et al. (2016) showed that far-IR colour temperature was also lower by ~ 20 per cent on average compared to dust temperatures estimated from a 4-point spectral energy distribution fit of the *Herschel* data. It is also worth noting that \bar{T}_{int} provides an upper limit to the temperature of compact sources as its calculation assumes optically thin emission and a spherically symmetric density profile that peaks at the location of the 70- μm bright protostar. Deviations from these assumptions would lead to lower mass-averaged temperatures. As a consequence, in the remaining of the analysis, the quoted temperatures are computed




Figure 4. Far-infrared colour temperature obtained from the ratio of the *Herschel* 160 μm to 250- μm fluxes, versus the internal temperature obtained from the *Herschel* 70- μm flux for each protostellar ArTéMiS source. The solid yellow line represents the median value of the $\bar{T}_{\text{int}}/T_{\text{col}}$ ratio. The grey shaded area shows the 16th to 84th percentile range.

using:

$$T_{\text{dust}} = 1.2(\pm 0.2) T_{\text{col}} \quad (5)$$

with the exception of the sources that have $T_{\text{col}} > \bar{T}_{\text{int}}$, for which we used \bar{T}_{int} . Using equation (5) allows us to compute dust temperatures consistently for all ArTéMiS sources, something that the use of \bar{T}_{int} would not allow us to do as it requires the detection of a 70- μm source. Finally, note that these temperatures are estimated on the scale of the *Herschel* 250- μm beam, i.e. 0.23 pc at 2.6-kpc distance, which is slightly more than twice larger than the ArTéMiS beam itself. According to equation (1), this can lead to a systematic underestimate of dust temperatures of ~ 30 per cent for protostellar sources. The impact of this important systematic uncertainty on temperature is discussed in Section 6.

5.2 Masses

The mass of each ArTéMiS source is estimated assuming optically thin dust emission, uniform dust properties (temperature and dust emissivity) along the line of sight, and uniform dust-to-gas mass ratio. With these assumptions, the source mass is given by:

$$M_{\text{gas}} = \frac{d^2 F_{\nu}}{R_{d2g} \kappa_{\nu} B_{\nu}(T_d)}, \quad (6)$$

where d is the distance to the source, F_{ν} is the source flux, R_{d2g} is the dust-to-gas mass ratio, κ_{ν} is the specific dust opacity at frequency ν , and $B_{\nu}(T_d)$ is the Planck function at the same frequency and dust temperature T_d . Here, we used $R_{d2g} = 0.01$ and $\kappa_{\lambda} = 7.35 \left(\frac{\lambda}{350\ \mu\text{m}} \right)^{-2} \text{ cm}^2/\text{g}$ (e.g. Hildebrand 1983; Könyves et al. 2015). Regarding distances, for each field we used the average distance of the individual clumps lying within them, with the exception of SDC326.796 + 0.386, which has been excluded from the rest of this study since it is much closer than all the other sources (see Table 1). Finally, regarding the dust temperature, we use T_{dust} as defined in Section 5.1.3. As far as uncertainties are concerned, we used 30 per cent, 15 per cent, and 20 per cent uncertainty for F_{ν} , d , and T_{dust} , respectively, that we propagated in equation (6) using Monte Carlo uncertainty propagation.




Figure 5. Mass versus dust temperature for all ArTéMiS sources. The temperatures are here estimated on the *Herschel* 250- μm beam size (i.e. 0.23 pc), while the ArTéMiS flux used to compute the mass is estimated on the ArTéMiS beam size (i.e. 0.1 pc). The symbols are coded by the sources’ internal luminosities. Sources with no 70- μm association from the Molinari et al. (2016) catalogue are represented as black symbols. Among these, those that do not display any local 70- μm or 8- μm peaks on the individual images presented in Appendix C are represented as filled circles. Those that do present a visually identified mid-infrared peak in these individual images have in addition a larger empty circular symbol. The dashed lines show the mass sensitivity limits at the two extreme distances of the sources in our sample. The blue dotted line gives half the thermal critical Bonnor Ebert mass for a core radius of 0.05 pc as a function of temperature.

The dendrogram analysis done here provides boundaries for every *leaf* identified in the ArTéMiS images. While we can use these to define the physical boundaries of compact sources, it is not clear if such an approach is the best. First, in some cases, especially for starless sources, these boundaries seem to encompass sub-structures that just fail to pass the detection criterion (i.e. local minimum to local maximum amplitude larger than $3\sigma_{\text{map}}$). Also, nearly all high-angular resolution ($\leq 1''$) observations of similar sources show sub-fragmentation (e.g. Louvet et al. 2019; Sanhueza et al. 2019; Svoboda et al. 2019) casting doubts on the true physical meaning of the identified ArTéMiS compact sources. Our approach here is more generic: we compute the mass within the ArTéMiS beam solid angle at the location of the peak flux density of every identified *leaf*. Because the sources analysed here are all within a very narrow range of distances (see Table 1), the proposed approach provides a measure of compact source masses within a comparable physical diameter of $\sim 0.10 \pm 0.01$ pc.

6 THE MASS–TEMPERATURE–LUMINOSITY DIAGRAM

6.1 The ArTéMiS view

As protostars evolve with time, the temperature, luminosity, and mass of their envelopes change. The accretion history of these protostellar envelopes will define what their tracks will be on a mass versus dust temperature diagram. Large statistical samples of protostellar sources within star-forming regions can therefore help constraining the accretion histories of these objects. In Fig. 5, we show the mass versus dust temperature diagram for all identified ArTéMiS sources with masses estimated using the temperatures

given by equation (5) and the ArTéMiS peak flux density. On the same figure, we have added the mass sensitivity limits for the minimum and maximum distances of our sample. One advantage of a mass versus dust temperature diagrams over a more standard mass versus luminosity one is that all sources, starless and protostellar, can easily be represented on it.

Fig. 5 displays a couple of important features. First, we notice the presence of warm ($T_{\text{dust}} > 30$ K) starless sources, which might seem surprising at first. However, these sources are all located in very specific environments, that is in the direct vicinity of some of the more luminous young stellar objects we have mapped. For instance, starless sources #14, 17, 18, and 20 in the SDC328 field have all dust temperatures larger than 30 K (including the two warmest ones displayed on Fig. 5 at 44 K and 55 K) and are all located within a radius of 0.6 pc of sources #13 and #19. These two sources have internal luminosities of $\sim 8500 L_{\odot}$ and $\sim 55000 L_{\odot}$, respectively. According to equation (1), sources with such luminosities can warm up dust up to 30 K within a radius of 0.3 pc and 0.6 pc. It is therefore unsurprising to find starless sources with temperatures in excess of 30 K. However, it is unclear if such sources are gravitationally bound and will form stars in the future. As a reference, a Bonnor–Ebert sphere of 0.05-pc radius and 40 K gas temperature has a critical mass of $\sim 4 M_{\odot}$ (Ebert 1955; Bonnor 1956). In Fig. 5, we added, as a blue-dotted line, the critical half-mass Bonnor–Ebert relationship for a core radius of 0.05 pc, $\frac{1}{2} M_{\text{BE}}^{\text{crit}} = 1 \times \left(\frac{T}{20\text{K}}\right) M_{\odot}$. Starless sources below that line are very likely to be unbound structures.

An even more important feature of Fig. 5 is the presence of massive protostellar sources with masses beyond $30 M_{\odot}$ and the absence of equally massive starless counterparts. This is in line with the early result by Motte et al. (2007) on the lack of massive pre-stellar cores in Cygnus. We also note that a luminosity gradient seems to run from the low-mass low-temperature corner to the high-mass high-temperature one. These trends, however, are very much subject to the relative temperature difference between starless and protostellar sources. As noted in Section 5.1.3, the flux and temperature measurements used to build Fig. 5 are inconsistent with each other since they are estimated on different spatial scales, i.e. 0.1 pc and 0.23 pc, respectively. Because the temperature profiles of starless and protostellar sources scales are different, this inconsistency could create artificial trends in a diagram such as that of Fig. 5. We attempt to correct for it using equation (1) for protostellar sources and assuming that starless sources are isothermal. The mass-averaged temperature correction factor for protostellar sources is given by $(0.23/0.1)^{1/3} = 1.32$ [from equation (1), with $\beta = 2$]. The temperature of starless sources are left unchanged. The resulting corrected temperature versus mass diagram is shown in Fig. 6. On this figure, we see that the trends observed in Fig. 5 (i.e. protostellar sources being more massive than starless ones, and the presence of a diagonal luminosity gradient) are mostly still present, albeit with slightly decreased significance. All data (temperature, mass, and luminosity) used to produce that figure is provided in Table 2.

The correction we made on the source temperatures relies on the fact that our starless/protostellar classification is robust. However, as mentioned in Section 4, ~ 43 per cent of the ArTéMiS sources that do not have a 70- μm association from the Molinari et al. (2016) catalogue seem to have a 70- μm and/or 8- μm emission peak when looking at the individual source images provided in Appendix C (in Figs 5 and 6, these sources are marked as black empty circular symbols with a smaller black-filled symbol in them). Also, when observed with ALMA at high-angular resolution, single-dish starless sources observed in high-mass star-forming regions systematically fragment into a set of low-mass protostellar cores




Figure 6. Same as Fig. 5 but temperatures and masses have been rescaled to a common spatial scale of 0.1pc.

(e.g. Svoboda et al. 2019). This shows that the classification of sources as starless based on single-dish continuum observations (e.g. with *Herschel*) should be viewed with caution in these regions. The net impact of wrongly classifying a protostellar source as starless would be to underestimate its temperature and therefore overestimate its mass. In other words, the trends mentioned earlier can be strengthened only by correcting for such misclassifications. This is particularly true if the handful of cold massive sources above $10 M_{\odot}$ would turn out to be protostellar (as Fig. 6 shows, it is likely to be the case for at least three of these sources).

Finally, we also note that the relationship provided by equation (5), even though established on protostellar sources only, has been applied to all sources, including starless ones. This seems to be the most appropriate approach since the ratio \bar{T}_{int} over T_{col} does not appear to be a function of the internal luminosity (see Fig. 4). However, for completeness, we do show in Appendix D a version of the mass versus temperature diagram in which we used $T_{\text{dust}} = T_{\text{col}}$ for the starless sources while applying the same correction factors for the protostellar sources.

Given the relatively low number of sources in our sample, the trends mentioned above are rather speculative. Nevertheless, it remains interesting to determine whether or not one can recover these trends with simple models that mimic both core-fed and clump-fed accretion scenarios.

6.2 Accretion models

Following Bontemps et al. (1996), André et al. (2008), and Duarte-Cabral et al. (2013), we built a simple accretion model that is aimed at reproducing the evolution of a protostellar core as the central protostar grows in mass. The set of equations that describes the mass growth of a protostar, and the parallel mass evolution of the core, is:

$$\frac{dm_*}{dt} = \dot{m}_* \quad (7)$$

$$\frac{dm_{\text{core}}}{dt} = -\dot{m}_* + \dot{m}_{\text{clump}} \quad (8)$$

$$\dot{m}_* = \epsilon_{\text{cs}} \frac{m_{\text{core}}}{\tau_{\text{core}}} \quad (9)$$

$$\dot{m}_{\text{clump}} = \epsilon_{\text{cc}} \frac{m_{\text{clump}}}{\tau_{\text{clump}}}, \quad (10)$$

where m_* is the mass of the protostar, m_{core} is the mass of the core, \dot{m}_* is the mass accretion rate of the protostar, \dot{m}_{clump} is the mass accretion rate of the core from the clump, τ_{core} is the characteristic star formation time-scale on core scale, τ_{clump} is the characteristic star formation time-scale on clump scale, ϵ_{cs} is the star formation efficiency from core to star (the fraction of the core mass that is being accreted on to the protostar), and finally ϵ_{cc} is the core formation efficiency from clump to core (the fraction of the clump mass that ends up in a core). In the context of this set of equations, core-fed scenarios differentiate themselves from clump-fed ones by having $\dot{m}_{\text{clump}} = 0$. This is the framework Duarte-Cabral et al. (2013) worked in. The clump-fed models, on the other hand, are presented here for the first time. In the following, we explore both type of scenarios.

Equations 7–10 provide a description of the mass evolution of both the protostar and the surrounding core. However, in order to produce a mass versus temperature diagram, one needs to compute, in parallel to the mass evolution, the evolution of the luminosity of the system. To do this, we used the protostellar evolutionary tracks from Hosokawa & Omukai (2009). These are well adapted to the formation of massive stars. These tracks provide, for a given mass accretion rate and given protostar mass, the total luminosity of the system that includes both accretion luminosity and stellar luminosity. At each time-step of our numerical integration of equations (7)–(10), we linearly interpolate the luminosity between the closest tracks. Finally, using equations (1), (2), and (5), one can then compute the theoretical equivalent of Fig. 6.

In the context of core-fed scenarios, cores refer to the fixed-mass reservoir of individual protostars. In nearby low-mass star-forming regions, these cores have typical sizes ranging from 0.01pc to 0.1pc (e.g. Könyves et al. 2015, 2020). These can be understood as the typical sizes of the gravitational potential well’s local minima, decoupled from their larger scale surroundings. In the context of clump-fed scenarios, these cores are located within a larger scale minimum defined by the presence of a surrounding parsec-scale clump that continuously feeds the cores with more mass. While equations (7)–(10) do not explicitly refer to any size-scale, the calculation of the mass-averaged temperature, equation (1), does require setting a characteristic core scale. Here, we are limited by the spatial resolution of the ArTéMiS observations, i.e. ~ 0.1 pc at the distance of the observed regions. Hence, in the following models, we use $R_{\text{core}} = 0.05$ pc.

Fig. 7 shows a set of models with $\dot{m}_{\text{clump}} = 0$ (effectively core-fed models), and for six different initial (pre-stellar) core masses, $m_{\text{core}}(t=0) = [5, 10, 20, 40, 80, 160] M_{\odot}$, all with a radius $R_{\text{core}} = 0.05$ pc. As suggested by Duarte-Cabral et al. (2013), we set $\tau_{\text{core}} = 3 \times 10^5$ yr for all sources. Note that the exact value used for this time-scale does not change the shape of the modelled tracks; a shorter time-scale would only make the evolution faster. We also set $\epsilon_{\text{cs}} = 0.3$, lower than the value of 0.5 used in Duarte-Cabral et al. (2013) to represent the fact that the modelled cores are larger. In essence, the tracks presented in Fig. 7 are identical to those presented in Fig. 5 of Duarte-Cabral et al. (2013) (albeit the slightly different set of parameter values). While these models cover a similar range of mass and temperature as the ArTéMiS sources, they require the existence of massive pre-stellar cores that should reside in the top left corner of the plot. For the tracks describing the evolution of the most massive stars [$m_{\text{core}}(t=0) = [80, 160] M_{\odot}$], such starless sources are not present in our ArTéMiS sample. But one could argue though that such core-fed models provide a good description of the data for initial core masses $m_{\text{core}} \leq 30 M_{\odot}$ which, according to the models, would form stars with $m_* \leq 9 M_{\odot}$. These same intermediate-mass




Figure 7. Core-fed models. Each track has been computed for a different initial core mass, from bottom to top $m_{\text{core}}(t=0) = [5, 10, 20, 40, 80, 160] M_{\odot}$. The coloured symbols represent the position of the cores at times $t = [9 \times 10^3, 9 \times 10^4, 1.8 \times 10^5, 2.7 \times 10^5, 3.6 \times 10^5]$ yr. The colour codes the stellar mass at these times as displayed by the colour bar. The background grey symbols are those presented in Fig. 6. Note that sources with $M_{\text{gas}} < \frac{1}{2} M_{\text{BE}}^{\text{crit}}$ have been removed.




Figure 8. Clump-fed models. Each track has been computed for a different clump mass, from bottom to top $m_{\text{clump}} = [100, 200, 400, 800, 1600, 3200] M_{\odot}$. The coloured symbols represent the position of the cores at times $t = [3 \times 10^4, 3 \times 10^5, 6 \times 10^5, 9 \times 10^5, 1.2 \times 10^6]$ yr. The colour codes the stellar mass at these times as displayed by the colour bar. The background grey symbols are those presented in Fig. 6. Note that sources with $M_{\text{gas}} < \frac{1}{2} M_{\text{BE}}^{\text{crit}}$ have been removed.

tracks also explain the presence of luminous objects (i.e. $L_{\text{int}} \geq 10^3 L_{\odot}$) with low associated core masses as sources that arrive at the end of their accretion phase.

Fig. 8 shows a set of tracks with $m_{\text{clump}} \neq 0$ (effectively clump-fed models) and $m_{\text{clump}} = [100, 200, 400, 800, 1600, 3200] M_{\odot}$. They all start with the same initial core mass $m_{\text{core}}(t=0) = 1 M_{\odot}$, the typical Jeans mass in dense molecular clumps. We also set $\epsilon_{\text{cc}} = 0.1$, $\epsilon_{\text{cs}} = 0.3$, and $\tau_{\text{clump}} = \tau_{\text{core}} = 1 \times 10^6$ yr, i.e. the clump crossing time controls the infall. This assumption remains valid as long as the time to regenerate the mass of the core, i.e. $\frac{m_{\text{core}}}{m_{\text{clump}}} \tau_{\text{clump}}$, is shorter than the core free fall time. This is verified at all times in

the models. We set a longer time-scale for clump-fed models than for core-fed models since the gas density of clumps is necessarily lower than that of the cores embedded within them. However, as for the core-fed models, the exact value used for the time-scale in the clump-fed models does not change the shape of the tracks. Finally, the core accretion phase is stopped once $t > \tau_{\text{clump}}$. Note that the point of this paper is not to proceed to a thorough examination of the parameter space of the proposed model but rather to evaluate if such models could generate a reasonable agreement with the observations. As we can see in Fig. 8, these models do also cover a similar range in mass and temperature as the observations and are able to explain the formation of the most massive stars without the need for massive starless sources. In addition, the modelled tracks evolved along the evolutionary gradients that we tentatively see in the observations. These models are therefore rather promising in the context of trying to pinpoint the physical mechanisms lying behind the mass accretion history of the most massive stars.

One could argue that the spatial resolution of the ArTéMiS data presented here (i.e. 0.1 pc) is not enough to probe individual pre/protostellar cores, and that the ArTéMiS sources are therefore likely to be sub-fragmented. While this might be true, it is also likely that the measured ArTéMiS flux of each source is dominated by the brightest unresolved core lying within the ArTéMiS beam. In fact, there is evidence that this is indeed the case as Csengeri et al. (2017) observed eight of the most massive ArTéMiS sources presented here with ALMA at $\sim 3''$ resolution, nearly three times better resolution than ArTéMiS and corresponding to a size scale of ~ 8000 AU. On that scale, the fraction of the ALMA flux locked in the brightest ALMA compact source is between 50 per cent and 90 per cent of the total flux. Also, Csengeri et al. (2018) presented ALMA observations of source SDC328#19 (one of the two warmest sources presented in, e.g. Fig. 5, at an angular resolution of $0.17''$, i.e. ~ 500 AU at 2.75 kpc). There, no sub-fragmentation is observed.

A comparison between our ArTéMiS observations and models on scales smaller than the ArTéMiS beam requires a set of extra assumptions and is therefore most uncertain. Such comparison is provided in Appendix E.

7 CONCLUSIONS

The key observational constraint regarding core-fed star formation is the existence of massive pre-stellar cores. The most massive starless sources identified here have masses of $\sim 30 M_{\odot}$ in a 0.1-pc source size, which is three to four times less massive than the most massive protostellar sources identified in the observed fields (within the same size). Taken at face value, this would suggest that the most massive ArTéMiS sources we identified keep growing in mass while simultaneously feeding massive protostar(s) at their centre, and that clump-fed models describe best the formation of massive stars. Our data though do not exclude the possibility of core-fed star formation for intermediate-mass stars. Therefore, a transition regime could exist between core-fed and clump-fed star formation scenarios around $m_* = 8 M_{\odot}$.

Most of the ArTéMiS sources studied here are likely to be sub-fragmented into a number of unresolved individual cores. A larger fragmentation level in our ArTéMiS protostellar sources, compared to the starless ones, could invalidate our former conclusion and instead favour core-fed scenarios. High-angular resolution observations on 1000 AU scale of massive 0.1-pc-sized sources, both starless and protostellar, have indeed revealed sub-fragmentation (e.g. Bontemps et al. 2010; Palau et al. 2013; Sanhueza et al. 2019; Svoboda et al. 2019). There is, however, no evidence that starless