Astronomy & Astrophysics manuscript no. main
March 7, 2022
©ESO 2022

arXiv:2203.02208v1 [astro-ph.GA] 4 Mar 2022

The 700 ks *Chandra* Spiderweb Field I: evidence for widespread nuclear activity in the Protocluster

P. Tozzi[1], L. Pentericci[2], R. Gilli[3], M. Pannella[4], F. Fiore[5], G. Miley[6], M. Nonino[5], H.J.A. Röttgering[6], V. Strazzullo[5], C. S. Anderson[7], S. Borgani[4,5,8,9], A. Calabrò[2], C. Carilli[10], H. Dannerbauer[11,12], L. Di Mascolo[4], C. Feruglio[5], R. Gobat[13], S. Jin[11], A. Liu[14], T. Mroczkowski[15], C. Norman[16,17], E. Rasia[5,8], P. Rosati[18], A. Saro[4,5,8,9]

[1] INAF - Osservatorio Astrofisico di Arcetri, Largo E. Fermi, I-50122 Firenze, Italy e-mail: paolo.tozzi@inaf.it
[2] INAF - Osservatorio Astronomico di Roma, Via Frascati 33, I-00040 Monteporzio (RM), Italy
[3] INAF - Osservatorio di Astrofisica e Scienza dello Spazio, via Piero Gobetti 93/3, 40129 Bologna, Italy
[4] Università di Trieste, Dipartimento di Fisica, Sezione di Astronomia, via Tiepolo 11, 34143 Trieste, Italy
[5] INAF-Osservatorio Astronomico Trieste, via Tiepolo 11, 34123 Trieste, Italy
[6] Leiden Observatory, PO Box 9513, 2300 RA Leiden, The Netherlands
[7] Jansky Fellow of the National Radio Astronomy Observatory, P. O. Box 0, Socorro, NM 87801, USA
[8] INFN - Sezione di Trieste, via Valerio 2 - 34127 Trieste, Italy
[9] Institute of Fundamental Physics of the Universe, via Beirut 2, 34151 Grignano, Trieste, Italy
[10] National Radio Astronomy Observatory, P. O. Box 0, Socorro, NM 87801, USA
[11] Instituto de Astrofísica de Canarias (IAC), 38205 La Laguna, Tenerife, Spain
[12] Universidad de La Laguna, Dpto. Astrofísica, 38206 La Laguna, Tenerife, Spain
[13] Instituto de Física, Pontificia Universidad Católica de Valparaíso, Casilla 4059, Valparaíso, Chile
[14] Max Planck Institute for Extraterrestrial Physics, Giessenbachstrasse 1, 85748 Garching, Germany
[15] European Southern Observatory (ESO), Karl-Schwarzschild-Str. 2, D-85748 Garching, Germany
[16] Space Telescope Science Institute, 3700 San Martin Dr., Baltimore, MD 21210, USA
[17] Johns Hopkins University, 3400 N. Charles Street, Baltimore, MD 21218, USA
[18] Dipartimento di Fisica e Scienze della Terra, Universitá degli Studi di Ferrara, via Saragat 1, I-44122 Ferrara, Italy

March 7, 2022

ABSTRACT

Aims. We present an analysis of the 700 ks *Chandra* ACIS-S observation of the field around the radio galaxy J1140-2629 (the Spiderweb Galaxy) at $z = 2.156$, focusing on the nuclear activity in the associated large-scale environment.
Methods. We identify unresolved X-ray sources in the field down to flux limits of 1.3×10^{-16} and 3.9×10^{-16} erg/s/cm^2 in the soft (0.5-2.0 keV) and hard (2-10 keV) band, respectively. We search for counterparts in the optical, NIR and submm catalogs available in the literature to identify X-ray sources belonging to the protocluster and derive their X-ray properties.
Results. We detect 107 X-ray unresolved sources within 5 arcmin (corresponding to 2.5 Mpc) of J1140-2629, among which 13 have optical counterparts with spectroscopic redshift $2.11 < z < 2.20$, and 1 source with photometric redshift consistent with this range. The X-ray emitting protocluster members are distributed approximately over a $\sim 3.2 \times 1.3$ Mpc^2 rectangular region. An X-ray spectral analysis for all the sources within the protocluster, shows that their intrinsic spectral slope is consistent with an average $\langle\Gamma\rangle \sim 1.84 \pm 0.04$. Excluding the Spiderweb Galaxy, the best-fit intrinsic absorption for 5 protocluster X-ray members is $N_H > 10^{23}$ cm^{-2}, while other 6 have upper limits of the order of few$\times 10^{22}$ cm^{-2}. Two sources can only be fitted with very flat $\Gamma \leq 1$, and are therefore considered Compton-thick candidates. The 0.5-10 keV rest frame luminosities of the 11 Compton-thin protocluster members, corrected for intrinsic absorption, are greater than 2×10^{43} erg/s. These values are typical for the bright end of a Seyfert-like distribution and significantly greater than X-ray luminosities expected from star formation activity. The X-ray luminosity function of AGN in the volume associated to the Spiderweb protocluster in the range $10^{43} < L_X < 10^{44.5}$ erg/s, is at least 10 times higher than that in the field at the same redshift and significantly flatter, implying an increasing excess at the bright end. The X-ray AGN fraction is measured to be $25.5 \pm 4.5\%$ of the spectroscopically confirmed members in the stellar mass range $\log(M_*/M_\odot) > 10.5$. This value corresponds to an enhancement factor of $6.0^{+9.0}_{-3.0}$ for the nuclear activity with $L_{0.5-10keV} > 4 \times 10^{43}$ erg/s with respect to the COSMOS field at comparable redshifts and stellar mass range.
Conclusions. We conclude that the galaxy population in the Spiderweb Protocluster is characterized by enhanced X-ray nuclear activity triggered by environmental effects on Mpc scales.

Key words. galaxies: clusters: general – galaxies: active – X-rays: galaxies: clusters

1. Introduction

Protoclusters are defined as overdense regions in the high-z Universe which are expected to evolve into massive, virialized clusters of galaxies at the present epoch (see Overzier 2016, for a review). Finding and characterizing protoclusters is key to study the large scale structure of the Universe and the transformational processes that affect star formation and nuclear activity in the member galaxies, such as powerful mergers, gas cooling, and feedback effects. In practice, although a protocluster is observationally identified as a high-z structure that is overdense compared to its surroundings, the dynamical state of the structure,

and therefore whether it will evolve into a low-z cluster, is usually highly uncertain (see, e.g. Muldrew et al. 2015). At present, there are no standard methods to systematically search for *bona fide* protoclusters, and this is the reason why biased-tracer techniques are often used to identify interesting targets. One of these consists in searching around high redshift, powerful radio galaxies, that are often found to be the beacon of overdense regions (see Miley et al. 2006). Independently of the selection method, in order to confirm their nature and to trace the activity of several processes occurring during their rapid evolution, protocluster candidates must be followed-up intensively with multiwavelength campaigns.

In this respect, X-ray observations play a key role, particularly for studying: (1) unresolved emission from active galactic nuclei (AGN) indicative of accretion onto nuclear supermassive black holes (SMBH), (2) the less intense emission from strongly star forming galaxies, and (3) diffuse emission from hot gas and/or relativistic plasma. Unfortunately, $z \gtrsim 2$ protoclusters are usually very faint in the X-rays and therefore only deep, high-resolution observations can provide useful information in this band. Until now only a few deep ($t_{\rm exp} \geq 200$ ks) observations of protoclusters have been made with *Chandra* or XMM-Newton. In a few cases a faint, extended emission has been interpreted as the signature of a low-mass virialized halo (Gobat et al. 2011; Valentino et al. 2016; Wang et al. 2016). In some other studies, the thermal or relativistic nature of the diffuse emission is still unclear (Gilli et al. 2019; Champagne et al. 2021). Despite the scanty X-ray data on protoclusters, a clear aspect is that to search and characterize the diffuse emission in high-z protocluster and, at the same time, the unresolved emission from AGN members, high sensitivity and angular resolution ($\sim$ 1 arcsec) are both required.

Regarding the X-ray nuclear activity, there is increasing evidence that this is enhanced in overdense, protocluster regions. Lehmer et al. (2009) and Digby-North et al. (2010) found, respectively, 12 and 4 X-ray sources with $L_{2-10keV} \geq 3-5 \times 10^{43}$ erg s^{-1} probably associated to the protoclusters identified in the SSA22 and HS1700+64 fields. These numbers exceed by a factor of 10-20 the X-ray source counts in deep fields at the same redshifts, although the uncertainties are large (see also Krishnan et al. 2017). The enhancement of X-ray AGN in high-z protoclusters is particularly impressive when compared with the overdensity of X-ray sources in evolved clusters at $z \sim 0.5$ that is only a factor of $\sim$ 3 (Martini et al. 2006; Branchesi et al. 2007; Kocevski et al. 2009; Ehlert et al. 2015). However, we note that the physical quantity that should be measured is not the X-ray source density, but the AGN fraction among protocluster members, hence the ratio of this fraction to the same quantity measured in the field, defined as the enhancement factor $f_{\rm enh}$. To be meaningful, this quantity should be evaluated in a well defined stellar mass range, and possibly differentiated between star-forming and passive galaxies, in order to compare the full distribution of AGN accretion rates in the protocluster members to those in field galaxies at comparable redshift. A robust characterization of $f_{\rm enh}$ in protoclusters that accounts for the dependence on redshift and on the mass scale is still not available. Furthermore, the lack of quantitative estimates on the luminosity function of AGN in protoclusters and on their obscuration implies large uncertainties on a related, important quantity such as the energy budget released in the surrounding medium by the nuclear activity.

Finally, another relevant missing piece of information in protoclusters is the measurement of the hot component in the surrounding diffuse baryons (the forming intracluster medium, ICM), that can be used to constrain the virialization status of the halo. A characterization of the proto-ICM, including its morphology, thermodynamical and chemical properties, can provide an independent and direct measure of the heating efficiency associated to the feedback and, at the same time, of the gas accretion into the protocluster potential well. In addition, feedback processes not only affect the ICM, but can also couple directly with the member galaxies. An interesting example was recently found in the field of the high redshift quasar SDSSJ1030 (observed by *Chandra* with a 500 ks exposure), where a protocluster at $z \sim 1.7$ centered on a bright FRII radio galaxy, shows star-forming member galaxies apparently triggered by the AGN radio jet (Gilli et al. 2019). If confirmed, this phenomenon may shed new light on the strong interplay between the member galaxies and the surrounding medium in the first stages of formation of large structures.

As of today, the available data are providing tantalizing evidence for the complex and intricate phenomena that occur in such high-density, rapidly evolving regions at cosmic noon. A comprehensive physical picture of the accretion, aggregation and phase transformation of baryons in protoclusters will require large sample statistics (still limited by the number of currently known protoclusters) and a multiwavelength approach, including high-resolution X–ray observations. Here we shall address these issues using deep *Chandra* ACIS-S observation ($\sim$ 700 ks) of the remarkable Spiderweb Galaxy (J1140-2629) complex at $z = 2.156$ (redshift obtained from low-resolution spectroscopy by van Ojik 1995). The central, powerful radio galaxy, embedded in a giant Lyα halo (Pentericci et al. 1997; Miley et al. 2006), is surrounded by a $\gtrsim$ 2 Mpc–sized overdensity (corresponding to a radius of 4 arcmin) of star forming galaxies (Lyα and Hα emitters), dusty starbursts, and red galaxies likely in the process of shaping a nascent red sequence (Zirm et al. 2008). The complex around the Spiderweb Galaxy is considered to be a typical protocluster region which is expected to evolve into a massive cluster in less than 1.5 Gyr, with the radio galaxy itself showing the properties of a cD progenitor (Miley et al. 2006).

In this paper we focus on the nuclear activity in the protocluster members, while the physics associated to the diffuse emission surrounding the Spiderweb Galaxy will be investigated in a companion paper based on X-ray and SZ data (Tozzi et al. in preparation; DiMascolo et al. in preparation). The paper is organized as follows. In Section 2 we summarize the properties of the Spiderweb Galaxy complex on the basis of previous observations. In Section 3 we briefly describe the multiwavelength data set used in this work. In Section 4 we outline the X-ray data acquisition and reduction techniques. Section 5 deals with the identification all the X-ray sources in the field and their number counts. In Section 6 we list the galaxies associated to the large scale structure (protocluster members) and characterize their nuclear activity in terms of X-ray luminosity and spectral properties. In Section 7 and 8 we present the X-ray luminosity function and the AGN fraction of the protocluster members, respectively. In Section 9 we discuss the enhancement factor implied by our new measurements and discuss our results in comparison to previous X-ray studies of protoclusters. Possible extensions of this work in promising directions are briefly discussed in Section 10. Finally, our conclusions are summarized in Section 11. For the sake of clarity, we refer to the radio galaxy and its immediate surroundings as the Spiderweb Galaxy, or J1140-2629, while "Spiderweb Complex" indicates the large scale structure potentially associated to the radio galaxy in a broad redshift range ($2.0 < z < 2.3$). The term "protocluster", is instead used to include only the galaxies in the narrower red-

shift range $2.11 < z < 2.20$ (see also Jin et al. 2021), assuming that all the protocluster members will end up in a massive ($\sim$few $\times 10^{14} M_{\odot}$) cluster today. Throughout this paper, we adopt the seven-year WMAP cosmology with $\Omega_{\Lambda} = 0.73$, $\Omega_m = 0.27$, and $H_0 = 70.4$ km s^{-1} Mpc^{-1} (Komatsu et al. 2011). In this cosmology, at $z = 2.156$, 1 arcsec corresponds to 8.473 kpc, the Universe is 3.13 Gyr old, and the lookback time is 77% of the age of the Universe. Quoted error bars correspond to a 1 σ confidence level, unless noted otherwise.

2. The Spiderweb Complex: previous observational campaigns and main results

Since its discovery in 1994, the Spiderweb galaxy (J1140-2629) and the Spiderweb Complex have become an intensively studied region. In the last 25 years, 54 papers have been published about this field including observations in radio, sub-mm, IR, optical and X–ray wavelengths. In this section we summarize the main results that have been obtained so far, focusing on the protocluster environment.

2.1. Discovery and peculiarities

The Spiderweb Galaxy was discovered during the implementation of an ESO Key Program designed to find high-z radio galaxies (HzRG) by targeting ultra-steep spectrum radio sources (Roettgering et al. 1994). The object was noticed as peculiar because of its unusually clumpy and bent radio morphology, coupled with an exceptionally high rotation measure ($\sim$ 6200 rad/m), indicating that the radio synchrotron jet is surrounded and its propagation constrained by a dense ($10^{-1} - 10^{-2}$cm^{-3}) external medium (Carilli et al. 1997; Pentericci et al. 1997; Athreya et al. 1998). Optical follow-up classified it as a narrow emission line galaxy at $z = 2.156$ (see Roettgering et al. 1997), revealed a clumpy optical continuum morphology, and a spectacular Lyα halo (still one of the largest known in the Universe) with a luminosity of 42.5×10^{44} erg/s elongated along the jet. The total stellar mass inferred from the K-band luminosity is $10^{12} M_{\odot}$ (Pentericci et al. 1997), confirmed by Spitzer data (Seymour et al. 2007), making the Spiderweb one of the most massive, high-z galaxies known. Using HST ACS observations, the clumpy features, embedded in the 200 kpc Lyα halo, whose distribution resembles that of flies trapped in a spiderweb, have been shown to correspond to $\sim$ 10 star-forming satellite galaxies moving with peculiar velocities of several hundred km/s, presumed to be merging with the central radio galaxy (Miley et al. 2006).

Eventually, Nesvadba et al. (2011) found that the optical and NIR spectrum of the nucleus of J1140-2629 shows a broad, spatially unresolved Hα line (blended with [NII]) having a FWHM of $\sim$ 14900 km/s. The presence of broad nuclear lines is rare for optical counterparts of high redshift radio sources, typically classified as TypeII AGN. The black hole mass is estimated to be $M_{\bullet} \sim 2 \times 10^{10} M_{\odot}$, putting J1140-2629 a factor of $\sim$ 2 above the local $M_{\bullet} - \sigma$ relation, in agreement with its expected evolution. Taken together, the observed characteristics suggest that the Spiderweb is indeed a massive radio galaxy forming at the center of a dynamically evolving protocluster region, that is likely to evolve into a brightest cluster galaxy. The Spiderweb is, therefore, one of the most revealing laboratories for studying the evolution of a protocluster in the early Universe, and may represent a typical phase in the formation of massive galaxy clusters with prominent central galaxies.

2.2. Identification of protocluster members and their properties

The search for protocluster members of the Spiderweb Complex was based on a series of works with spectroscopic follow-up and narrow band imaging in the field. A first sample of 50 candidate Lyα emitters in a 6×6 arcmin^2 field (Kurk et al. 2000), provided support to the existence of several star forming galaxies in the protocluster. Out of these candidates, 15 were eventually confirmed to have redshift in the range 2.16 ± 0.02 with VLT-FORS spectroscopy by Pentericci et al. (2000). One of these confirmed sources shows clear signs of AGN activity with very broad Lyα and CIV emission lines, while all the others show just narrow Lyα emission. More extensive VLT imaging in NB0.38, B, R and I bands with FORS2 and in the J_s, H, K_S and NB2.07 bands with ISAAC, identified 40 Hα candidate emitters and 44 extremely red objects (EROs) consistent with $z \sim 2$, whose density increases towards the central galaxy suggesting a physical association (Kurk et al. 2004b). Estimated star formation rates (SFRs), without dust correction, are in the range 2-50 $M_{\odot}$/yr and 0.4-4.3 $M_{\odot}$/yr for the Hα and Lyα emitter candidates, respectively. The SFR per unit mass, estimated from K-band magnitudes, are 2.5 times higher than in local massive clusters.

In a spectroscopic follow-up in the IR band with ISAAC on the VLT, additional 9 Hα emitter candidates within 0.6 Mpc from J1140-2629 have been confirmed as protocluster members (Kurk et al. 2004a). The narrowness of the emission lines of 8 out of the 9 confirmed sources indicates that they are powered by star formation dominated by very young (< 100 Myr) stellar population with moderately high metallicity. Only in one of the emitters the Hα appears extremely broad with a FWHM exceeding 5000 km/s. Their contribution to the SFR density in the protocluster shows it to be at least an order of magnitude larger than that found at $\langle z \rangle = 2.23$ in the HDF-N.

Near-infrared spectroscopy of narrow-band selected Hα emitters with MOIRCS on Subaru presented in Shimakawa et al. (2014, 2018) significantly increased the number of spectroscopically confirmed protocluster members. An additional 4 protocluster members were identified among 13 Hα emitter candidates using ALMA CO(3–2) (ν_{rest} = 345.796 GHz) observations (Tadaki et al. 2019), all of them also detected in CO(4–3) by Emonts et al. (2018). Recently, Jin et al. (2021) obtained 46 CO(1-0) detections in the range $2.09 < z < 2.22$, using 13 individual pointings with ATCA.

Photometric studies also helped to characterize the protocluster population. IR Surveys indicated an overdensity of 3-4 σ with respect to the field at 24μm (Mayo et al. 2012), with Spitzer IRAC colors (Galametz et al. 2012), and with Herschel SPIRE (Valtchanov et al. 2013; Rigby et al. 2014). The distribution of IR protocluster galaxy candidates confirms the approximate filamentary shape of the protocluster, previously obtained using optical narrow band selection. In the submm band, an overdensity of a factor of 4 is observed with APEX LABOCA at 870 μm (Dannerbauer et al. 2014). The submm flux distribution at the protocluster redshift would imply a SFR larger by about an order of magnitude with respect to the Hα derived values. The sample of narrow-band selected candidates was extended to 68 Hα emitters with the MAHALO Deep Cluster Survey (Shimakawa et al. 2018) with the Subaru Telescope, 17 of which are newly discovered. In addition, $Bz'K_s$ selection identified 34 distant red galaxies, half of which have photo-z consistent with being protocluster members.

Overall, the galaxy population of the Spiderweb Complex appears to be a mix of active and star forming galaxies, and

already massive, passive galaxies that constitute a nascent red sequence (Kodama et al. 2007; Zirm et al. 2008, 2012), and that approximately follow and elongated structure extending ∼ 5 Mpc in diameter. Tanaka et al. (2013) found that the weak red sequence is consistent with a formation redshift in the range $3 < z < 4$. In addition, Koyama et al. (2013) found that the properties of the Hα emitters are consistent with them having formed a large part of their stellar mass at $z > 2$, but observed in a still intense star-forming phase. The presence of both star forming and red galaxies may be related to a typical phase associated to environmental effects driven by the protocluster collapse, that accelerates their formation and evolution.

Finally, we note that four out of six (60%) massive H_α emitters (HAEs) ($M_* > 10^{11} M_\odot$) have been already identified as bright X-ray sources (see Pentericci et al. 2002; Croft et al. 2005). Also, Croft et al. (2005) confirmed two X-ray sources as protocluster members using LRIS Keck spectroscopy, and classified both as AGN due to the presence of NV, CIV and [CIII] in the spectrum, bringing the total number of X-ray emitting members to five. This finding suggests that protocluster members that host X-ray AGNs may be in the transition stage from dusty star-forming galaxies to passive populations.

2.3. Role of the new X-ray observations

The nature of the galaxy population of the Spiderweb Protocluster will be further investigated in this work thanks to the addition of our deep X-ray data. Our study relies on a collection of 252 unique sources in the field that have spectroscopic or narrow-band/photometric redshift in the range $2.0 < z < 2.3$ (see Section 3). A detailed list of all the spectroscopic members and color or narrow-band selected candidates, and the identification of the Spiderweb Complex candidates are shown in Section 3.

Using our deep *Chandra* observation we shall explore the following aspects: i) the frequency of X-ray nuclear activity among the protocluster members; ii) the enhancement of nuclear activity in the protocluster; iii) the distribution of phases in the diffuse baryons in the halo of the Spiderweb galaxy, and the relation with its strong nuclear activity; iv) the presence of diffuse thermal emission possibly associated to the virialization of a central halo. In this paper we focus on the AGN population. The presence of hot, diffuse baryons around the Spiderweb Galaxy and its relation to its nuclear activity will be discussed in two papers based on X-ray and SZ signal (Tozzi et al. in preparation; Di Mascolo et al. in preparation). The main results and the data products can be found on the project webpage[1].

3. Multiwavelength data used in this work

In this work we shall exploit part of the large set of multi-wavelength data publicly available for the Spiderweb Complex field. We use the HST images for the central regions and the Subaru images, which cover the entire *Chandra* FOV, to identify the optical counterparts of the X-ray sources. In particular, we use a 23 ks exposure in the F814W band with ACS (HST Proposal 10327, PI H. Ford), and B, r', z' exposures obtained with the Suprime camera at the Subaru telescope and covering a $\approx 30 \times 30$ arcmin^2 field. We also use VLT-VIMOS imaging data in the U band, and VLT-HAWKI images in Y, H and K bands. The Suprime B, r', and z' raw data were retrieved from the SMOKA archive (Baba et al. 2002). The images have been reduced with a custom pipeline due to the 4 ports readout layout of the camera chips. Images have been *overscan*, *master bias* and *master flats* corrected. We then used an improved version of own pipeline as described in (Nonino et al. 2009), which currently uses Scamp (Bertin 2006) with GAIA-EDR3 as astrometric reference, and Swarp (Bertin et al. 2002) for the final coaddition. A similar approach has also been used for the VLT-VIMOS and VLT-HAWKI images, with the raw data retrieved from the ESO Archive. The instrumental signatures removal has been performed using IRAF[2]. For the NIR HAWKI images the appropriate darks were retrieved from the ESO Archive, stacked and applied to the raw data. For the optical data the bias and the flats have been used to create static weights which account for chips defects. For the NIR data darks and flats has been instead used.

We use optical and NIR photometry to estimate host galaxy masses and rest-frame luminosities of the X-ray AGNs in our sample. A detailed description of the photometric measurements, multi-wavelength catalogue, photometric redshift and stellar mass estimation will be given elsewhere (Pannella et al., in preparation). In the following, for the sake of completeness we briefly summarise the data and procedures relevant here.

VLT and Subaru stacked images have been used to create a multi-wavelength catalogue with 7 passbands (U,B,r',z',Y,H,Ks) from U to K. In table 1 we summarise the main properties of dataset used. We use SExtractor in dual image mode, to measure photometry. The K_S-band HAWK-I image has been adopted as the primary detection image, because it represents the best compromise, among all available bands, between the need for a robust tracer of galaxy stellar mass and a good angular resolution (≃0.4"), which simplifies catalog assembly and photometry measurements. The whole catalog contains 2,279 objects over the K_S image field of approximately 84 arcmin^2 and down to an AB magnitude of 22.8 (i.e., the image 5σ limiting magnitude). The images used here have very different resolutions. Rather than convolving all images to the worse resolution, which would result in a significant loss of information, we account for this in the estimate of aperture colors by applying PSF-matching corrections based on the growth-curve of point-like sources. To limit uncertainties in such corrections we use 2" diameter apertures to sample the galaxy spectral energy distributions (SEDs).

Photometric redshifts are computed using EAZY (Brammer et al. 2008). Global photometric zero points are adjusted iteratively by minimizing the difference between photo-zs and the available spectroscopic redshifts (e.g. Gabasch et al. 2004; Ilbert et al. 2010; Pannella et al. 2015). By comparison with the available spectroscopic sample, the rms of $(z_{phot} - z_{spec})/(1 + z_{spec})$ is ∼6%. Stellar masses are derived using a modified version[3] of FAST (Kriek et al. 2009), adopting Salpeter (1955) IMF, Bruzual & Charlot (2003) stellar population synthesis models, delayed exponentially declining star formation histories (SFHs), the Calzetti et al. (2000) dust extinction law (with $A_V = 0$ to 4), and solar metallicity. In deriving the stellar mass, we use the photo-z obtained using the FAST code, while we use the spectroscopic redshifts when available.

To compile a comprehensive list of spectroscopic redshifts, we use a collection of catalogs of all the spectroscopically confirmed protocluster members, member candidates selected on the basis of color or narrow-band photometry, and random sources with redshift in the field. The 17 catalogs, based on 12 different

[1] http://www.arcetri.inaf.it/spiderweb/

[2] IRAF was distributed by the National Optical Astronomy Observatory, which was managed by the Association of Universities for Research in Astronomy (AURA) under a cooperative agreement with the National Science Foundation.

[3] https://github.com/cschreib/fastpp

Table 1. Optical and NIR Data.

Filter	Instrument /Telescope	FWHM (arcsec)	5σ (AB)	Science Archive /Project ID(s)
U	VIMOS/VLT	0.86	26.4	ESO - 383.A-0891
B	S-Cam/Subaru	1.12	25.8	SMOKA - o17409
r'	S-Cam/Subaru	0.73	25.2	SMOKA - o17409
z'	S-Cam/Subaru	0.65	24.4	SMOKA - o10144
Y	HAWK-I/VLT	0.46	24.5	ESO - 088.A-0754, 091.A-0106
H	HAWK-I/VLT	0.58	23.2	ESO - 088.A-0754, 091.A-0106
K_S	HAWK-I/VLT	0.40	22.8	ESO - 088.A-0754, 091.A-0106

Notes. The first to fourth columns indicate filter name, instrument/telescope, seeing FWHM, and 5σ limiting magnitude in a 2 arcsec diameter aperture, respectively. In the fifth column we list the science archive and proposal ID of the raw dataset.

publications, are listed in Table 2. The catalogs have a significant overlap, since several of them include and extend previous results. The total number of unique sources with any redshift information in the field is 252, out of which 161 have spectroscopic redshifts. We also compile a complementary list of 91 protocluster member candidates, identified thanks to narrow-band or color selection in previous works. Most (90%) of the sources (both confirmed and candidate members) are included in a box of 6.3×3.7 arcmin^2 (corresponding to 3.13×1.84 Mpc at $z = 2.156$) roughly centered on the Spiderweb galaxy and elongated along the E-W direction. Only 9 sources with redshift information are outside a circle of 5 arcmin centered on the Spiderweb Galaxy, and none of them are in the range $2.0 < z < 2.3$.

The distribution of the 161 available spectroscopic redshifts is shown in Figure 1. Clearly the majority of the redshift values are close to the Spiderweb redshift $z = 2.156$, since protocluster member candidates have been mostly targeted. We consider a redshift interval $\Delta z = 0.3$ roughly symmetric with respect to the Spiderweb redshift, corresponding to a velocity difference of $\Delta v \sim c \times \Delta z/(1+z)$ associated to the Hubble flow. This interval spans all the sources possibly associated to the Spiderweb large scale structure. If we consider the range $2.0 < z < 2.3$, corresponding to a maximum $\Delta v \sim 28500$ km/s, we find 112 sources. We do not expect to see a normal distribution for the redshift, therefore we adopt the biweight estimator for the central value and the standard deviation, as suggested by Beers et al. (1990) for sample size $n \geq 100$[4]. We find $z_{BI} = 2.15754$ and $\sigma_{BI} = 0.0208612$. Therefore, two and three sigma intervals centered on z_{BI} are $2.116 < z < 2.1996$ and $2.095 < z < 2.220$, respectively. We decide to assume the 2σ intervals (approximated as $2.11 < z < 2.20$) as the protocluster redshift range.

In the range $2.11 < z < 2.20$, corresponding to a maximum $\Delta v \sim 8500$ km/s, we find 96 sources. We compute the formal velocity dispersion σ_v in this range and find $\sigma_v \sim 1490 \pm 150$ km/s. This value should be compared to that expected for a flat distribution in the same redshift range as expected for sources following the Hubble flow, which is ~ 2700 km/s. We conclude that the sources are clearly detached from the Hubble flow, and show a global velocity dispersions that would correspond to a very high mass ($M > 10^{15}\,M_\odot$), virialized single halo. Clearly, given the mass scale and the distribution of sources in the sky, the virialized, single halo hypotesis is not realistic. In addition, we notice that excluding the new spectroscopic sources identified by Jin et al. (2021) with the CO(1-0) line, the velocity dispersion drops to $\sigma_v = 980 \pm 100$ km/s (that would correspond to a temperature of ~ 6 keV for a virialized halo). As already shown in Jin et al. (2021), this points out that the CO sources have a different dynamical behaviour, and are probably distributed along a large scale filament or a galaxy superprotocluster. In addition, half of the CO emitters have measured velocities larger than the escape velocities of any realistic virialized halo hosting the Spiderweb Galaxy, suggesting that the outer regions traced by the CO sources are still far from being accreted onto the main halo. In this work we do not attempt to investigate further the dynamical structure of the Spiderweb complex. However, the number of spectroscopically confirmed members in the Spiderweb Complex is now the highest for a single structure at $z > 2$ and a detailed, spatially resolved dynamical analysis will be a relevant piece of information to constrain the current status and the future evolution of the structure.

In summary, in this paper we consider the 96 sources included in the narrow redshift bin $2.11 < z < 2.20$ as belonging to the protocluster, while the additional 16 sources included in the range $2.0 < z < 2.3$ are considered part of the large scale structure associated to the Spiderweb Complex, and most likely are not dynamically bound. The corresponding redshift intervals are shown in Figure 1 with vertical dashed lines.

4. X-ray data reduction

The Spiderweb Galaxy was observed with a Chandra Large Program observation of 700 ks with ACIS-S granted in Cycle 20 (PI P. Tozzi). The observations were completed in the period November 2019 - August 2020, splitted in 21 separate Obsid. To this data set we add the first X-ray observation with ACIS-S, dating back to June 2000 for a total of 39.5 ks. All the 22 Obsid used in this work are listed in Table 3.

Data reduction is performed starting from the level = 1 event files with `CIAO 4.13`, with the latest release of the *Chandra* Calibration Database at the time of writing (`CALDB 4.9.4`). We run the task `destreak` to flag and remove spurious events (with moderate to small pulse heights) along single rows, which is particularly important for ACIS-S4. We run the tool `acis_detect_afterglow` to flag residual charge from cosmic rays in CCD pixels. Finally, since all our new observations are taken in the VFAINT mode we run the task `acis_process_events` with the parameter `apply_cti=yes` to flag background events that are most likely associated to cosmic rays and reject them. With this procedure, the ACIS particle background can be significantly reduced compared to the standard grade selection. The data is then filtered to include only the standard event grades 0, 2, 3, 4, and 6. The *level 2* event files generated in this way are visually inspected for flickering pixels or hot columns left after the standard reduction but we find none after filtering out bad events. We also carefully inspect the image of the removed photons, particularly to verify whether

[4] See equations 5, 6 and 9 in Beers et al. (1990)

Table 2. Catalogs of spectroscopically confirmed protocluster members and member candidates used in this work.

Reference	Selection method	Members
Spectroscopic		
Pentericci et al. (2000)	Lyα emission	15
Kurk et al. (2004a)	Hα emission	9
Croft et al. (2005)	Lyα emission	7 [17]
Doherty et al. (2010)	Hα emission	2 [2]
Kuiper et al. (2011)	OII emission	15 [1]
Tanaka et al. (2013)	OII/OIII emission	4 [10]
Dannerbauer et al. (2014)	Specz	11 [4]
Shimakawa et al. (2014)	Hα emission	23
Tadaki et al. (2019)	CO emission	4
Jin et al. (2021)	CO emission	46
Color or narrow-band selected		
Pentericci et al. (2002)	photo-z	3
Kurk et al. (2004b)	Lyα emitter	40
Kurk et al. (2004b)	Hα emitter	40
Kurk et al. (2004b)	EROs	44
Dannerbauer et al. (2014)	FIR photo-z	2 [4]
Shimakawa et al. (2018)	Hα emitter	68
Tadaki et al. (2019)	Hα emitter	9

Notes. In the second column we list the identification/selection method, while in the third column we report the number of confirmed or candidate members of the Spiderweb complex ($2.0 < z < 2.3$). Occasionally, the number of random sources in the Spiderweb field outside the range $2.0 < z < 2.3$ is reported in square brackets. The catalogs have been obtained independently in different works, but often overlap significantly since are often derived from the same datasets. The total number of unique sources is 252. We find 161 sources with spectroscopic redshifts, out of which 112 in the range $2.0 < z < 2.3$.

Table 3. *Chandra* observations of the Spiderweb Galaxy Complex.

Obsid	date	Exptime (s)
898	2000-06-06	27 378
21483	2019-11-18	49 312
22924	2020-02-26	33 475
22923	2020-02-28	34 598
23175	2020-03-01	29 938
22925	2020-03-06	39 541
22926	2020-03-07	43 375
22921	2020-03-17	24 642
23190	2020-03-17	24 736
23069	2020-03-23	19 648
23201	2020-03-26	27 613
21482	2020-04-02	34 492
22927	2020-04-02	32 534
23205	2020-04-03	23 514
22928	2020-04-10	29 572
23215	2020-04-13	19 803
21484	2020-04-22	46 504
22929	2020-04-23	51 231
23186	2020-06-13	14 770
22922	2020-07-22	18 487
21481	2020-08-13	40 383
22905	2020-08-04	49 291

Notes. In the last column we list the exposure time after data reduction.



Fig. 1. Upper panel: Distribution of the 161 spectroscopic redshifts in the Spiderweb field. Lower panel: redshift distribution in the range $2.0 < z < 2.3$. Vertical dashed lines show the redshift of the Spiderweb Galaxy as measured with low resolution spectroscopy by van Ojik (1995) (black line), the range adopted to define the Spiderweb Complex (magenta lines) and the protocluster (green lines), as discussed in the text.

we have pile-up effects[5]. We find that slightly more than 100 net counts in the total band are removed at the position of the nucleus of J1140-2629 across the 22 exposures. This indicates that the source, which is by far the brightest in the field, suffers some amount of pile-up. However, we will verify *a posteriori* that this effect does not impact the spectral analisis of the nucleus and therefore it is not worth to give up the VFAINT cleaning to recover less than 1% of the flux in the nucleus. In particular, the VFAINT cleaning is very important for the characterization of the diffuse emission, which is presented in detail in further papers from the collaboration, focused on the Inverse Compton emission associated to the radio jets (Carilli & Anderson 2022; Anderson & Carilli 2022) and on the thermal component (Tozzi et al. in preparation; see also Di Mascolo et al. in preparation).

Finally, time intervals with high background are filtered by performing a 3σ clipping of the background level. The light curves are extracted in the 2.3–7.3 keV band, and binned with a time interval of 200 s. The time intervals when the background

[5] https://cxc.cfa.harvard.edu/ciao/ahelp/acis_pileup.html

exceeds by 3σ the average value, are removed with the tool `deflate`. The final total exposure time after data reduction and excluding the dead-time correction amounts to 715 ks (corresponding to the LIVETIME keyword in the header of *Chandra* fits files), including the first observation. The average removed time interval for the new observations amounts to only $\sim$ 0.3% of the observing time, except for Obsid 898, where $\sim$ 30% of the exposure time has been removed due to high background time intervals. The monochromatic exposure maps are created at the energies of 1.5 and 4.5 keV for the soft and hard band respectively, by running the tools `mkinstmap` and `mkexpmap`.

The 22 *level 2* event files are merged together with the tool `reproject_obs`, using the reference coordinates of Obsid 21483. The exposure maps are reprojected onto the reference frame of Obsid 21483 with the tool `reproject_image`, and then are added together weighting each map by the fractional exposure time of the corresponding Obsid. Images for source detection are created in the 0.5-2 keV, 2-7 keV and total (0.5-7 keV) bands, with no binning to preserve the full angular resolution of $\sim$ 1 arcsec (FWHM) at the aimpoint (1 pixel corresponds to 0.492 arcsec). The images include all the CCD that were "on" while observing, namely ACIS-S3 (where the aimpoint is always located), ACIS-S2, ACIS-S4, and ACIS-I3 (except Obsid=898, in which also ACIS-S5 and ACIS-I2 were on). Since the aimpoint is approximately the same for the 22 Obsid, and considering the geometry of the CCD positions, the effective exposure is optimal within a radius of few arcmin from the J1140-2629 position, while it is significantly lower in a wide annulus covered by the flanking CCDs. We identify a region with a radius of 5 arcmin centered on J1140-2629, roughly coincident with all the points with an effective area larger than 240 cm^2. This radius corresponds to $\sim$ 2.5 Mpc at $z = 2.156$. Within this region the sensitivity falls by a factor of $\sim$ 2 from the center to the border. Beyond this radius, the flanking CCDs cover a larger area up to a radius of $\sim$ 14 arcmin, but with a much lower sensitivity and a significantly degraded angular resolution. The two regions drawn on the monochromatic 1.5 keV exposure map of the field are shown in Figure 2. The region at off-axis angle between 5 and 14 arcmin is not studied in this work.

In Figure 3, upper panels, we show an image of the central part of the field (a box 12×9 arcmin by side), in the soft and hard bands. The images have been smoothed with a $\sim$ 1 arcsec kernel to emphasize the unresolved sources. The color scales are the same in both bands (max 200 counts in log scale) and therefore it is possible to appreciate the larger background in the 2-7 keV band with respect to the soft 0.5 -2 keV band. In the lower panels, close-up images of the Spiderweb galaxy (a box of 3×2 arcmin by side, no smoothing applied) are shown in the soft and hard bands. The radio contours in the 10 GHz band obtained with JVLA (Carilli & Anderson 2022) are shown in red. From a visual inspection, it is possible to note that the unresolved sources surrounding the Spiderweb Galaxy are distributed approximately along the jet directions, as already noticed for the spatial distribution of the narrow-band selected sources. Finally, we note that the extended emission in the soft band appears not only along the jets but also in the direction perpendicular to the jet axis.

To perform spectral analysis, we extract the spectra and compute the ancillary response file (ARF) and redistribution matrix file (RMF) at the source position for each observation separately with the command `mkarf` and `mkacisrmf`. Our default spectral analysis of unresolved sources uses a local background, that, given the small extent of the sources is directly extracted from a source-free region on the same chip. This applies both to the unresolved sources and the diffuse emission. A more careful treatment will be devoted to the analysis of the nucleus of the Spiderweb galaxy, which is embedded in the diffuse emission. For most of the sources in the field, the soft and hard band fluxes are estimated simply by converting the count rate obtained by aperture photometry to energy flux through the conversion factors. We prefer aperture photometry to a more model-dependent photometry relying on the PSF modelization, since this last procedure can introduce systematic biasing effects that may be hard to control. On the other hand, the Poissonian noise associate to aperture photometry can be handled with simple statistical methods. The fluxes obtained in this way account for the full uncertainty due to Poisson noise, allowing us to obtain a reliable measurements for the entire source population across 3 orders of magnitude. The derivation of the conversion factors is detailed in the following subsections.



Fig. 2. Time-weighted, monochromatic exposure map at 1.5 keV of the Spiderweb Complex field. The maximum value is 470 cm^2. The innermost (red) circle shows the circular region with radius of 5 arcmin ($\sim$ 2.5 Mpc at $z = 2.156$) that is investigated in this work. All the points inside this circle have an effective area larger than 240 cm^2. The outer annulus, with radii of 5 and 14 arcmin, has a much lower effective exposure, and a degraded angular resolution, and it is not included in the present analysis.

5. X-ray sources in the Spiderweb Complex

In this paper we focus on the field of view within a radius of 5 arcmin (corresponding to $\sim$ 2.5 Mpc at $z = 2.156$) centered on the nucleus of the Spiderweb, which results in a total of 78.5 arcmin^2. This region includes the solid angle covered by our observation with a sensitivity larger than about 1/2 of the largest sensitivity reached at the aimpoint, and with an angular resolution better than 3 arcsec (FWHM). This allows us to have a smooth sky coverage, therefore avoiding large corrections when evaluating the number counts.



Fig. 3. Upper panels: *Chandra* X-ray images of the Spiderweb Galaxy field in the soft 0.5-2 keV (left) and hard 2-7 keV band (right). A box of 12×9 arcmin by side centered on J1140-2629 is shown. The images have been smoothed with a $\sim$ 1 arcsec Gaussian kernel to emphasize the unresolved sources. Sources at the outskirts of the image appear larger due to the effect of the increase of the PSF. Lower panels: close-up of the Spiderweb Galaxy (3×2 arcmin by side) at full angular resolution, in the soft (0.5-2 keV, left) and hard (2-7 keV, right) band. Extended emission is clearly visible in both bands. Red contours show radio emission observed in the 10 GHz band with the JVLA (see Carilli & Anderson 2022) at levels of 0.03, 0.2, 2 and 20 mJy/beam.

5.1. X-ray source detection

X-ray source detection is performed with an hybrid procedure combining visual inspection and a standard detection algorithm (`wavdetect`). The goal is to maximize the completeness of the final source list. In the first step, we consider the entire FOV included in the $\sim$ 14 arcmin radius, and visually identify 238 sources in at least one of the three images (soft, hard and total band), with approximate position and size.

Then, we run `wavdetect`[6] setting the scale parameter to `scales = "2 4 8 16 32"` and the detection threshold set to `sigthresh = 1e-05`. This parameter should be approximately equal to the number of pixels in the image, a choice that implies about one background fluctuation detected as source. Since we are searching for sources within 5 arcmin, we have about 1.17×10^6 pixels, setting `sigthresh = 1e-05` means approximately 12 strong background fluctuations detected as sources[7]. Our choice, therefore, is optimal for selection of unresolved sources down to low signal and low significance, with an acceptable level of contamination. Considering that our final selection will be based on a threshold in the S/N of aperture photometry, in our final list sources will show a S/N significantly higher than the lower limit of the `wavdetect` selection, implying also a much lower number of spurious sources with respect to the initial selection. We expect to have a completeness very close to unity, particularly in the inner 5 arcmin, where the angular resolution is a few arcsecs. We note, however, that running `wavdetect` with pixel scales below and equal to 32 is less efficient for detecting extended sources, or sources at off-axis angle $\sim$ 7 arcmin and larger, where the angular resolution of *Chandra* rapidly deteriorates. Since we focus on the sky region at off-axis angle < 5 arcmin, this choice is not affecting our results. In addition, a visual

[6] https://cxc.cfa.harvard.edu/ciao/threads/wavdetect/

[7] https://cxc.cfa.harvard.edu/ciao/ahelp/wavdetect.html#plist.sigthresh

inspection confirms that there are no relevant extended sources in the field within 5 arcmin. The extended emission surrounding the Spiderweb Galaxy, instead, will be discussed separately.

We find 249, 289 and 381 unresolved source candidates in the soft, hard and total-band images, respectively, selected by `wavdetect`. We match the `wavdetect` detected sources with the visually selected sample assuming a matching radius of 1 arcsec, and collect the `wavdetect` positions for all the matched sources, with the exception of the source candidates identified only by visual inspection. Then, we obtain a master catalog with 398 sources. All the visually-identified sources are recovered by the `wavdetect` algorithm but 11, all of which are at more than 5 arcmin from the Spiderweb, and therefore have no impact on this work. We define a circular extraction region for each source to perform aperture photometry. The extraction radius $r_{\rm ext}$ is computed with a simple relation depending only on the off-axis angle θ. We set $r_{\rm ext} = 2.55\times(0.67791-0.0405083*\theta+0.0535066*\theta^2)$ arcsec (see Rosati et al. 2002, and references therein). We find that this choice ensures that at least 95% of the expected flux is included in the circular extraction region, both in the soft and hard band[8]. We manually adjust position and size of the extraction regions to avoid overlapping of nearby source candidates: in three cases we manually reduce $r_{\rm ext}$ by one pixel (0.5 arcsec) and shift the centroid by 1 arcsec; in three other cases we simply reduce the $r_{\rm ext}$ by three pixels (1.5 arcsec). We verify *a posteriori* that, on average, the photometry of the six sources (flagged in Table A) change less than 10%, with a negligible impact on our final results. The background is estimated locally in an annulus around each source (outer radius $r_{\rm ext}+3''$, inner radius $r_{\rm ext}+1''$) after removing nearby unresolved sources. This choice, thanks to the depth of the total exposure, ensures a fair sampling of the background, and takes into account local variation of the background.

We estimate the Poisson probability of being a background fluctuation Pi for each candidate source i following the procedure described in Puccetti et al. (2020). First, we compute a radius $r_{\rm max}$ (with $r_{\rm max} < r_{\rm ext}$) within which the S/N is maximum. Then, we compute

$$P_i = e^{-B_i}\Sigma_{j=C_i,\infty}(B_i)^j/j! \quad (1)$$

where B_i is the estimated background counts within $r_{\rm max}$, and C_i is the total counts within $r_{\rm max}$. Puccetti et al. (2009) showed that assuming $P_i < 2.0\times10^{-5}$ ensures a very high reliability, so that we can use this threshold to select a sample of sources with a negligible spurious component. On the other hand, computing the completeness as a function of P_i requires extensive simulations to take into account the sensitivity and the PSF effects across the field of view. Therefore, for the completeness we prefer to use a criterion directly relying on the S/N from aperture photometry. This approach has been shown to be reliable in deep *Chandra* survey such as the Chandra Deep Field South, where we adopted a threshold of $S/N > 2.0$ to compute the point-source sky coverage (see Giacconi et al. 2001; Tozzi et al. 2001). In this work we combine these two criteria to exploit at best the sensitivity of the *Chandra* data keeping at the same time a very high purity of the sample.

We split our candidate list according to the distance from the Spiderweb Galaxy, finding 264 candidates at more than 5 arcmin and 134 detection candidates within 5 arcmin. For the remaining of the paper, we consider only the candidate detections in the central FOV. We compute P_i in the soft, hard and total (0.5-7 keV) bands for each source, and consider a detection any source that satisfy $P_i < 2.0\times10^{-5}$ in any of the three bands. This step selects 107 source candidates. Then, we apply our completeness criterion $S/N > 2.0$, and find that 9 out of 107 sources do not reach the S/N threshold in any band. However, we find that 5 of them have $S/N > 1.8$ in the hard band, and after a careful revision by visual inspection, we decide to keep them in the final source list, while discarding the remaining 4 sources.

In addition, we search for robust candidates that may have been missed by our reliability criterion, by searching for sources with $P_i > 2\times10^{-5}$ in all the three bands, but $S/N > 2$ in at least the soft or hard band. We find only 4 of them and all have a P_i value very close to our threshold (of the order of $3-4\times10^{-5}$). After a visual inspection, we decide to include them in the final sample, which, after this final step, includes 107 sources.

Incidentally, the same selection procedure identifies a final list of 176 sources in the outer ring, that has a field of view 6.8 times larger than the central 5 arcmin radius circle. This shows that the detected source density of in the outer ring is on average 4 times lower. Considering that sources at more than 5 arcmin from the aimpoint have significantly larger errors on position and photometry due to the rapid degradation of the PSF, and that the multiwalength data coverage essentially drops at distance > 5 arcmin, we strengthen the conclusion that the data quality in the field beyond 5 arcmin does not allow us to further extend the study of the Spiderweb Complex with the current dataset. The analysis of the field at distances larger than 2.5 Mpc from the Spiderweb is potentially interesting, in particular to investigate the possible extension of the Spiderweb Complex at radii larger than 2.5 physical Mpc, but it requires additional data and must be postponed to a future work.

5.2. Measurement of source fluxes

The exposure-corrected count rates are estimated by rescaling the measured net count rate by the ratio of the exposure map values at the ampoint to the emission-weighted exposure within the extraction region. The average monochromatic exposure maps values at the aimpoint are 470.8 cm^2 and 367.4 cm^2 in the soft and hard band, respectively. Soft- and hard-band fluxes are computed from the net count rates corrected for vignetting and assuming the typical conversion factors $C_{\rm soft}$ and $C_{\rm hard}$ in the corresponding band. The conversion factors clearly depends on the spectral shape, that is well known to depend on average on the source flux (see Tozzi et al. 2001), and it can be approximately described with a power law with a slope in the range $1.2 < \Gamma < 1.9$. Slopes in the range $\Gamma = 1.7-1.9$ are typical of unabsorbed AGN, as measured in deep surveys like the Chandra Deep Field South (CDFS, see Liu et al. 2017, and references therein). Harder spectra are mostly due to intrinsic absorption, which depends on the intrinsic column density and source redshift. Clearly, this information is not available for the full catalog. We proceed by assuming an average redshift of 1.5 (representative of AGN in deep X-ray surveys like the CDFS, see Figure 23 in Luo et al. 2017), and a typical intrinsic absorption $N_H \sim 10^{22}$ cm^{-2}. We find that the conversion factor obtained with these assumptions, and allowing for at least a factor of 3 variation in the average redshift and absorption, are well included in the range that we obtain assuming $1.2 < \Gamma < 1.9$. Therefore, without making a detailed description of the spectral shape of our unresolved sources, we adopt the full range $1.2 < \Gamma < 1.9$ and consider the corresponding variation in the conversion factors as a systematic uncertainty. This is a fair choice for the estimation of the source flux and a reliable mea-

[8] For details see `https://space.mit.edu/cxc/marx/tests/PSF.html`

Table 4. Conversion factors

Count rate band (keV)	Energy band (keV)	Conversion factor cgs/counts/s
0.5 - 2.0	0.5 - 2.0	$(8.77 \pm 0.65) \times 10^{-12}$
2.0 - 7.0	2.0 - 7.0	$(1.87 \pm 0.11) \times 10^{-11}$
2.0 - 7.0	2.0 - 10.0	$(2.66 \pm 0.35) \times 10^{-11}$

Notes. The conversion factors are computed at the aimpoint and corrected for Galactic absorption. The central value and the associated error correspond to a power law spectrum with slope in the range $1.2 < \Gamma < 1.9$, as discussed in the text.

surement of the number counts. The X-ray sources, identified as members of the Spiderweb Complex, will be further investigated with a detailed spectral analysis.

After correcting for a Galactic absorption column density of $N_{\rm H} = 3.18 \times 10^{20}$ cm^{-2}, we obtain $C_{\rm soft} = (8.77 \pm 0.65) \times 10^{-12}$ cgs/ counts/s in the soft band. In the hard band, we derive two conversion factors of $C_{\rm hard} = (1.87 \pm 0.11) \times 10^{-11}$ cgs/counts/s and $C_{\rm hard} = (2.66 \pm 0.35) \times 10^{-11}$ cgs/count/s to convert from the 2-7 keV count rate to the 2-7 keV and 2-10 keV fluxes, respectively. Note that we apply a factor 1.05 to account for the average flux loss outside the extraction region. Note also that while the net detected counts in the hard band refer to the 2–7 keV energy range, we compute the energy fluxes in the 2–10 keV band, for a better comparison with the literature, despite our unresolved sources showing very little signal above 7 keV. The list of 107 sources with X-ray position, net count rate in both bands and soft and hard fluxes is shown in Table A. A negative count rate may be measured when a source is detected only in one band, in which case we report the 2-σ upper limit for the energy flux.

5.3. X-ray number counts

To derive the number counts, we first compute the point-source sky coverage as a function of the energy flux. To do that, we first create images of the soft and hard background. These are obtained by removing all the detected sources and replacing the signal in the extraction region with a Poisson realization of the emission measured in the background annulus. Then, using the same conversion factors, and the monochromatic exposure maps, we compute the solid angle in which a source of a given soft or hard flux is detected according to our completeness criterion $S/N > 2.0$. As shown in Figure 4, the sky coverage rapidly drops to zero at fluxes of $1.3{\times}10^{-16}$ erg/s/cm^2 and $3.9{\times}10^{-16}$ erg/s/cm^2 in the soft and hard band, respectively[9]. The deepest X-ray field to date (and in the future until a new X-ray mission with arcsec resolution will be operating), is provided by the CDFS (Giacconi et al. 2001; Rosati et al. 2002; Luo et al. 2017) which has reached the unparalleled depth of 7 Ms. The comparison with the sky coverage of the CDFS shows that the field of view analyzed in this work is 5 times smaller than the maximum solid angle covered in the CDFS. The flux limit is about 13 and 10 times higher than in the CDFS in the soft and hard band, respectively. We note that the flux limits of the two fields roughly scale as the ratio of the total exposure times (a factor of 10), despite the sensitivity loss due to the contamination build-up over ACIS. The reason is that we take advantage of the larger effective area



Fig. 4. Thick lines show the point-source sky coverage of the innermost region of 5 arcmin of the Spiderweb field investigated in this work, in the soft (red) and hard (blue) band. Thin lines show the point-source sky coverage in the CDFS from Luo et al. (2017).

of the back-illuminated CCD in ACIS-S with respect to the front illuminated CCDs in ACIS-I (used for the CDFS), partially compensating the sensitivity loss.

The cumulative number counts (logNlogS) are computed directly as

$$N(>S) = \Sigma_{S_i > S} \frac{1}{\Omega_i}, \quad (2)$$

where S_i and Ω_i are the flux and the corresponding sky coverage of the $i^{\rm th}$ source, respectively. To compute the soft and hard number counts, we consider only the sources with $S/N > 2$ in the corresponding band. We select 72 sources in the soft and 96 sources in the hard band, that, combined together, add up to a total of 107 unique sources within the 5 arcmin radius.

The soft band logNlogS is shown in Figure 5. The shaded area show the formal 1σ uncertainty, obtained as the sum in quadrature of the statistical error due to the source statistics and the error on the source fluxes. The last error is, in turn, the sum in quadrature of the Poissonian photometric error and the systematic error associated to the conversion factors. The uncertainty is dominated by the source statistics at large fluxes, and by the photometric errors at low fluxes. In the same Figure we also plot the soft band logNlogS measured in the CDFS by Luo et al. (2017), and the predicted logNlogS based on the X-ray background synthesis model developed by Gilli et al. (2007)[10], that includes an exponential decline in the AGN space density at redshifts above $z = 2.7$ to cope with the results from wide-area surveys (e.g. SDSS Richards et al. 2006; Fan 2006). Since the Gilli et al. (2007) model includes only AGN, we add the steep galaxy number counts as measured by Luo et al. (2017) in the CDFS (see also Marchesi et al. 2020). Normal (star forming) galaxies are relevant only at fluxes well below 10^{-16} erg/s/cm^2, and therefore have a negligible impact on our results. In particular, the expected contribution of normal galaxies equals that of AGN at 10^{-17} erg s^{-1} cm^{-2} both in the 0.5-2 keV and 2-7 keV bands (see Figure 31 in Luo et al. 2017). The hard band logNlogS is shown in Figure 6. In both cases we find a very good

[9] Formally, the flux limit is defined as the flux at which the sky coverage drops to 1/10 of the full field of view of 78.5 arcmin^2.

[10] http://cxb.oas.inaf.it/, see also Hasinger et al. (2005).



Fig. 5. Cumulative number counts in the soft band in the Spiderweb field, within 5 arcmin from the Spiderweb Galaxy (red shaded area). The shaded area shows the uncertainty due to the combination of source statistics, photometric errors and systematic uncertainty on the conversion factors. The blue, dashed line shows the soft counts in the CDFS from Luo et al. (2017). The black solid line shows the prediction from the AGN model of Gilli et al. (2007) plus the contribution from normal galaxies from Luo et al. (2017).



Fig. 6. Cumulative number counts in the hard band in the Spiderweb field, within 5 arcmin from the Spiderweb Galaxy (red shaded area). The shaded area shows the uncertainty due to the combination of source statistics, photometric errors and systematic uncertainty on the conversion factors. The blue, dashed line shows the hard counts in the CDFS from Luo et al. (2017). The black solid line shows the prediction from the AGN model of Gilli et al. (2007) plus the contribution from normal galaxies from Luo et al. (2017).

agreement with the model expectation, while both model and counts in the Spiderweb Field appear to be $\sim$ 20% higher than the CDFS in the soft band at fluxes $\sim 10^{-15}$ erg/s/cm^2. The measured counts in the hard band show a $\sim$ 30% excess in a small range around fluxes $\sim 10^{-14}$ erg/s/cm^2 with respect to the model and CDFS. Overall, the logNlogS in the Spiderweb Field appear to be broadly in line with the expectations, with a small hint of excess on a limited flux range. Whether this excess is related to the presence of the Spiderweb Complex, will be investigated in Section 6.

6. Identification of X-ray emitting protocluster members

On the basis of the shallow *Chandra* observation (Obsid 898 Pentericci et al. 2002), only five X-ray sources have been identified among the spectroscopically confirmed or narrow-band selected protocluster members. These sources are firmly identified in the X-ray band as bright AGN in the 2–10 keV luminosity range 0.8-4$\times 10^{44}$ erg s^{-1}. No emission has been detected from any of the remaining galaxies in the protocluster down to a rest–frame 2–10 keV luminosity limit of 4×10^{43} erg s^{-1}. Four out of these five sources are aligned along the radio and X-ray diffuse emission (Croft et al. 2005). This is not surprising however, since confirmed and candidate protocluster members, irrespective of their selection, appear to be aligned in the same direction of the radio jets and of the X-ray emission. The relatively high AGN fraction estimated around the Spiderweb Galaxy, despite the low statistics, suggests that the AGN were probably triggered by the on-going formation of the protocluster. Another possible mechanism may be associated to the radio-mode feedback of the central radio galaxy, in a scenario similar to the observation of the protocluster in the SDSSJ1030 field (Gilli et al. 2019). However, the extension of the jets in the case of the Spiderweb Galaxy is limited to less than 100 kpc from the nucleus, making it a different case with respect to the SDSSJ1030 protocluster, where the jet extension is significantly larger (about 700 kpc between the two lobes). In any case, to demonstrate a trigger effect on the protocluster members, it is necessary to compute the fraction of X-ray emitting (at the Seyfert level or higher) members over the entire protocluster population. In this section we proceed to identify all the secure X-ray emitting protocluster members, with the final goal to estimate the AGN fraction in the Spiderweb system.

6.1. Matching X-ray sources to spectroscopic members

We match the X-ray sources with the optical source list, both with spectroscopic and photometric redshift, or narrow-band selected sources. Based on the accuracy of the X-ray centroid and of the error on the position of the CO selected sources, and neglecting the errors on the optical and NIR positions, we assume an initial matching radius of 1 arcsec for optical and NIR sources, and 2 arcsec for CO sources. Based on this values, we can immediately compute the expected number of false matches. Considering the entire FOV (a circle of 5 arcmin radius), and a cross section of π arcsec^2, we have a total fractional cross section of 2.8×10^{-3} for the 252 unique sources (including spectroscopic redshifts and narrow-band selected members). If the 107 X-ray sources are randomly distributed in this field of view, the expected number of random matches to protocluster members (confirmed or candidate) is 0.30. Therefore we expect significantly less than 1 spurious counterpart.

We obtain a list of 14 X-ray emitting protocluster members (including the Spiderweb), 13 of which matched with members with spectroscopic redshift (including the Spiderweb), and 1 (XID 73) with an ERO from Kurk et al. (2004b). Incidentally, we note than none of the *flies* is X-ray emitting. This is not surprising, since, given the measured mass range $10^8 < M_* < 10^{10} M_\odot$, and SFR range $0.5 - 26\, M_\odot$ yr^{-1} (Hatch et al. 2009), the corresponding cumulative X-ray emission from star formation only, is estimated to be more than an order of magnitude lower than our current flux limit. The selected sources are visually checked on

the X-ray, HST and Subaru images, to identify potentially spurious counterparts. We found that all the counterparts are unique within 1 arcsec except XID 90, that has a secondary counterpart that has also been selected as an ERO candidate at $z \sim 2.16$ Kurk et al. (2004b). Therefore, XID 90 may be considered, at worst, another X-ray member candidate without spectroscopic confirmation, similarly to XID 73.

The scatter plot of the offsets between all the X-ray counterparts and all the optical matches, from the catalogs we used in this work, is shown in Figure 7. The closest counterparts for each single source are marked with a white dot inside the red (spectroscopic) or green (photometric) point, showing that all the counterparts are well within 1 arcsec from the X-ray centroid, with most of them within 0.5 arcsec. We inspected the two sources (XID 12 and XID 80) that have an offset of $\sim$ 1 arcsec, and find that their X-ray centroid is not coincident with the X-ray peak, but is slightly displaced because of an apparent extension of the X-ray emission. The signal is too weak to classify these X-ray sources as extended [11], however, when considering the X-ray peak instead of the X-ray centroid, the offsets drop below 0.5 arcsec. Therefore, we find that, overall, we can safely assume that the offsets between the X-ray sources and the spectroscopic counterparts can be considered equal or less to 0.5 arcsec, that brings the expected number of spurious match below 0.1. We carefully inspected all the counterparts in the HST and Subaru images, finding no alternative optical counterpart candidates except those already mentioned. In the rest of the paper we will consider a sample of 12 X-ray spectroscopically confirmed members (excluding the Spiderweb) and an additional X-ray color-selected member candidate. Considering the 112 and 96 spectroscopically confirmed sources in the range $2.0 < z < 2.3$ and $2.11 < z < 2.20$, this corresponds to a fraction of $11.0 \pm 0.5\%$ and $13.0 \pm 0.5\%$, respectively. This fraction, however, does not take into account the dependence on the stellar mass, that is a key quantity that must be cosidered when computing the AGN fraction. This aspect will be discussed in Section 8.

In Figure 8 we show the X-ray image of the field in the 0.5-7 keV band with the X-ray emitting protocluster members marked with blue circles, and all the spectroscopically confirmed protocluster members marked with red circles. For a better visualization, here we show only a box of 460×240 arcsec entirely comprised within 5 arcmin from the Spiderweb galaxy (the FOV we consider), that includes all the spectroscopic protocluster members but two (non X-ray emitting). The filamentary distribution appears to be more evident in the X-ray sources than in the spectroscopically confirmed members, however, given the low statistics, the difference is not significant.

In Figure 9 we plot the azymuthally-averaged fraction of sources of a given class as a function of the distance from the Spiderweb Galaxy, finding that the protocluster members are clearly more clustered than the bulk of the X-ray sources, but also showing that there is no significant difference between the X-ray protocluster members and the spectroscopically-confirmed protocluster members. This is not surprising since the clustered distribution of the spectroscopic members has been already noticed in all the works listed in Table 2. This suggests that, in the current data, there is no evidence of enhanced X-ray activity closer to the Spiderweb Galaxy, if not for a marginally higher concentration of the X-ray emitting members. We checked whether the 15 sources considered to be part of the Spiderweb Complex but not in the protocluster (therefore belonging to the range $2.00 < z < 2.11$ and $2.20 < z < 2.30$) show a different behaviour, but we find their distribution similar to that of the protocluster members. We note that our conclusion on the X-ray activity being independent on the distance from the protocluster center, relies on the assumption of spectroscopic completeness, an aspect that, however, cannot be properly quantified given the composite nature of our spectroscopic sample.

[11] The extension may be associated to radio jets, an occurrence that will be tested when the wide-field, high-resolution radio images will be available.



Fig. 7. Scatter plot of the offsets for the 13 X-ray sources identified as members of the Spiderweb Complex, excluding the Spiderweb Galaxy. Red points corresponds to spectroscopic counterparts, while green points to color-selected counterparts. The offsets of the closest counterpart for each source are shown with a white dot inside the red or green points, while other points corresponds to other counterparts listed in independent catalog from Table 2. The large red circles correspond to 0.5 and 1 arcsec offsets.

Finally, in Figure 10 we plot the redshift distribution of the X-ray sources identified as protocluster members including the ERO (magenta filled histogram) compared to the redshift distribution of all the sources in the field with spectroscopic redshift in the range $2.0 < z < 2.3$. The two distributions are statistically equivalent, showing again that the X-ray emitting members are drawn uniformly in redshift among the spectroscopically confirmed members.

6.2. A note on the X-ray sources not associated to the protocluster

In addition to the X-ray sources identified as cluster members in the innermost 5 arcmin (13 spectroscopically confirmed, including the Spiderweb Galaxy, plus 1 color-selected members), we find counterparts for 9 sources with redshift $z < 2$, therefore in the foreground (6 of them are identified in the spectroscopic survey of X-ray sources in the field by Croft et al. (2005), and 3 are found in NED by chance). In the innermost 5 arcmin, we have 84 X-ray sources that have no redshift information. Clearly, the more important questions is whether we should expect more X-ray emitting protocluster members among this set of sources. To



Fig. 8. Hard (0.5-7.0 keV) band image of the center of the field (a box of 460 × 240 arcsec) with the X-ray emitting protocluster members marked with blue circles, and all the other spectroscopically confirmed protocluster members ($2.0 < z < 2.3$) marked with red circles. The image has been smoothed with a 1 arcsec Gaussian filter for clarity. X-ray members are labeled with the source ID, the original selection: Ha, Lya, ERO and CO for H_α, Ly_α, red color and CO-line selection, respectively, and spec for spectroscopic selection. Finally, the sources labeled with "X" have been identified as X-ray sources in the literature before this work, as described in Section 2.2.



Fig. 9. The fraction of sources within a given radius from the Spiderweb Galaxy. X-ray cluster members (shown in blue) have the same distribution of all the spectroscopically-confirmed protocluster members in the $2.11 < z < 2.20$ range (shown in red). On the other hand, the distribution of all the X-ray sources (dominated by the field) is clearly less clustered. The thin black line shows a perfectly uniform source distribution. The shaded area corresponds to an uncertainty of 1σ.



Fig. 10. Redshift distribution of the X-ray sources identified as protocluster members, including the one without spectroscopic confirmation (magenta filled histogram) compared to the redshift distribution of all the sources in the field with spectroscopic redshift in the range $2.0 < z < 2.3$. The dashed, vertical lines correspond to the protocluster redshift range.

give a first-cut estimate, we report here some preliminary results on the photo-z measurements of K_S band detected sources in the field (Pannella et al. in preparation), obtained with the method described in Section 3. Driven by the comparison of spec and photo-z in the field, we conservatively consider all the sources with $1.7 < z_{phot} < 2.6$ as potentially associated to the protocluster. We have 20 additional X-ray emitting sources in this sample. If we assume a flat distribution of the true redshift in this range, we expect ∼ 2 additional X-ray sources belonging to the protocluster ($2.11 < z < 2.20$).

Also, there are 58 X-ray sources within the central 5 arcmin that are matched with K_S sources with $z_{phot} < 1.7$ or $z_{phot} > 2.6$. Since it is very unlikely for these sources to have redshift in the protocluster range, we consider them as foreground and background sources. These leave out 15 X-ray sources without any redshift information. By visual inspection, most of these sources do not have possible counterpart in the HST and Subaru images, and therefore some of them may be high-z sources. Therefore, a reasonable assumption is to consider that all the 15 sources without any redshift information are outside the $2.11 < z < 2.20$ range. We also note that the fraction of X-ray sources without optical counterparts ($\sim$ 10%) is larger than that found in deep multiwavelength surveys like the CDFS (few %). This effect is not significant and depends on the depth and solid angle of the optical and NIR available coverage. This aspect will be further investigated in forthcoming papers (Pannella et al. in preparation).

As a final step, we select all the K_S band detected sources, complementary to the spectroscopic and narrow-band selected candidates we considered in this work, and find 380 sources within 5 arcmin with $1.7 < z_{phot} < 2.6$. Following the previous argument, we may expect additional $\sim$ 38 protocluster members without X-ray emission. If we focus on sources with $\log(M_*/M_\odot) > 10.5$, we may expect about 75% of these sources above this threshold (see Section 8), therefore additional 28 protocluster members. To summarize, preliminary results on the SED of galaxies detected in the Spiderweb field show that further spectroscopic follow up campaigns may increase the number of protocluster members by a $\sim$ 20−30%. Nevertheless, this simple, order-of-magnitude estimate of the X-ray protocluster members that can have been missed, is useful to bracket the AGN fraction and AGN enhancement factor in the Spiderweb protocluster, as discussed in Section 8 and Section 9.

6.3. Number counts in the $2.11 < z < 2.20$ redshift range

To quantify the overdensity of X-ray sources with respect to the field, we compute the X-ray number counts separately in a narrow redshift bin centered on the Spiderweb Galaxy. We decide to adopt the range $2.11 < z < 2.20$, and compute the logNlogS inside and outside this redshift range, both in the data and in the model of Gilli et al. (2007).

In Figure 11, left panel, we show the logNlogS in the soft band after excluding the X-ray sources identified in the $2.11 < z < 2.20$ range, compared to the model expectations. We find a substantial agreement, with the counts in the Spiderweb field being slightly below the model, in line with the CDFS data. In the right panel of Figure 11, we plot, instead, the soft-band logNlogS only for the redshift interval $2.11 < z < 2.20$ compared with the expected number counts from the model (black solid line). A much higher normalization is observed, and a different slope, showing that the excess is at relatively bright fluxes. Clearly the excess above fluxes 10^{-14} at soft erg/s/cm^2 is merely due to the presence of the Spidwerweb Galaxy. However, at fluxes $\sim 10^{-15}$ erg/s/cm^2 the excess can be robustly estimated and it turns out to be more than a factor of $\sim$ 5, and more than an order of magnitude at $\sim 10^{-14}$ erg/s/cm^2. We find the same behaviour in the hard band, as shown in Figure 12. Also in this case the excess is more than a factor of $\sim$ 5 at fluxes $\sim 10^{-15}$ erg/s/cm^2, and up to $\sim$ 10 at $\sim 10^{-14}$ erg/s/cm^2. Finally, we note that the strikingly different slopes in the range $2.11 < z < 2.20$ is due to the presence of relatively bright ($\sim 10^{-15}$ erg s^{-1} cm^{-2}) sources, thus are not affected by any means by our completeness correction. Therefore, we are able to confirm an excess in the number of X-ray sources per square degree at the same level estimated from spectroscopic, narrow-band, or color selected sources. The different shape of the number counts with respect to the model may be a hint that the X-ray population in this redshift range is not uniformly extracted from the field population, but there may be some environmental effect at play.

6.4. Properties of X-ray emitting protocluster members

How different is the population of X-ray protocluster members with respect to the bulk of the field X-ray sources in the Spiderweb Galaxy field? As a first check we plot the differential distribution of soft and hard-band fluxes in Figure 13. We note that in the soft and hard bands the distribution of X-ray fluxes of the protocluster members are slightly shifted towards higher values with respect to the entire X-ray sample.

Then, we proceed with a standard X-ray spectral analysis of the sources. We adopt a simple model consisting in an intrinsically absorbed power law, using the model components `zwabs` and `powerlaw` within Xspec. Galactic absorption is described with the model `tbabs`, and its value is fixed to 3.18×10^{20} cm^{-2} (HI4PI Collaboration et al. 2016). We also add a Gaussian line with intrinsic width below the spectral resolution, at an energy corresponding to the redshift neutral iron K_α line at 6.4 keV rest-frame, allowed to vary in energy by 5% around this interval (corresponding to a redshift uncertainty of $\Delta z \sim 0.1$). We formally measure the equivalent width in the observed band of this Gaussian component, without making an attempt to quantify the confidence level of a line detection, given the low signal regime. We leave the two shape parameter N_H and Γ both free to vary when a source has more than 40 counts in at least one observed band (0.5-2 keV or 2-7 keV), while we fix $\Gamma = 1.8$ in the other cases. The best fit values are shown in Table 5. In addition, we perform a simplified spectral analysis for the remaining 6 sources, where the intrinsic spectral slope is frozen to $\Gamma = 1.8$, and the intrinsic absorption is estimated. In the Appendix B we show the spectra of the 8 sources suitable for spectral analysis, along with the best fit model, and the spectra of the fainter 6 sources, despite the coarse binning. In this section we discuss the X-ray properties of the 13 protocluster members, while the Spiderweb Galaxy, except for the few basic spectral parameters, will be discussed in a companion paper (Tozzi et al. in preparation).

First, we note that we are able to measure the intrinsic slope with good accuracy in 8 sources, finding an average $\langle\Gamma\rangle = 1.75$. Source XID 86 has un unusual hard slope (with an hardness ratio $HR \sim 0$) and a measured equivalent width for the neutral Fe K_α line of $\sim$ 0.5 keV, close to the value expected for reflection-dominated, Compton-thick sources (see, e.g. Norman et al. 2002; Gilli et al. 2011; Iwasawa et al. 2020). We consider it a Compton Thick candidate with $N_H > 1.5 \times 10^{24}$ cm^{-2}. Therefore, the slope of its spectrum is not representative of the spectral shape of the intrinsic emission. If we exclude this source, we measure and average intrinsic slope of $\langle\Gamma\rangle = 1.86 \pm 0.05$, in line with expectations (see, e.g. Tozzi et al. 2006; Iwasawa et al. 2020). Another source with an unusually flat spectrum is XID 34. This source is the faintest among the protocluster X-ray members (17 net counts in the 0.5-7 keV band) and it cannot be meaningfully fitted with a frozen $\Gamma = 1.8$. We tentatively considered it another Compton-thick candidate.

For all the remaining sources, we are able to directly measure the intrinsic absorption or an upper limit for it. In Figure 14 we show the distribution of the intrinsic absorption for the 14 protocluster members (including the Spiderweb Galaxy). We also include upper limits (shown with the dashed histogram),



Fig. 11. Left panel: the soft band logNlogS in the Spiderweb field (within a radius of 5 arcmin) after excluding the X-ray sources identified in the $2.11 < z < 2.20$ range, compared to the expectations from the model of Gilli et al. (2007) in the same redshift range. The shaded area corresponds to an uncertainty of 1σ. Right panel: the soft band logNlogS in the Spiderweb field only for the redshift interval $2.11 < z < 2.20$, compared to the expectations from the model of Gilli et al. (2007) in the same redshift range. The shaded area corresponds to an uncertainty of 1σ. Note the different scales between the two panels.



Fig. 12. Left panel: the hard band logNlogS in the Spiderweb field (within a radius of 5 arcmin) after excluding the X-ray sources identified in the $2.11 < z < 2.20$ range, compared to the expectations from the model of Gilli et al. (2007) in the same redshift range. The shaded area corresponds to an uncertainty of 1σ. Right panel: the hard band logNlogS in the Spiderweb field only for the redshift interval $2.11 < z < 2.20$, compared to the expectations from the model of Gilli et al. (2007) in the same redshift range. The shaded area corresponds to an uncertainty of 1σ. Note the different scales between the two panels.

while we exclude the two Compton-thick candidates. The five sources (not including the Spiderweb) for which we robustly measure N_H, not surprisingly have $N_H > 10^{23}$ cm^{-2}, which is simply a consequence of the high-redshift and the low-signal regime. We also show 1σ upper limits for 6 sources, finding values $N_{\rm H} <$ few $\times\, 10^{22}$ cm^{-2}. The limited statistics and the low signal hamper any general conclusion on the distribution of $N_{\rm H}$ among the protocluster members. A visual comparison with the distribution of $N_{\rm H}$ expected in the model by Gilli et al. (2007) in the redshift bin $2.0 < z < 2.2$ (black line), normalized to the number of sources in our small sample, suggests that the number of strongly absorbed sources (7 sources with $N_{\rm H} > 10^{23}$ cm^{-2}) is broadly consistent with that expected in the field at the same redshift. We note, however, that the Compton-thin source with the highest intrinsic absorption that we are able to measure (XID 7, $N_{\rm H} = 4.0 \times 10^{23}$ cm^{-2}) is still > 4 times less absorbed than the most absorbed sources detected in protoclusters (see Gilli et al. 2019; Vito et al. 2020).

Having the intrinsic absorption, and an estimate of the intrinsic power law, we can compute the unabsorbed luminosity in the rest frame 0.5-2 and 2-10 keV bands. The unabsorbed luminosities of all sources but the two Compton-thick candidates and the Spiderweb Galaxy, are shown in Figure 15. All sources have a rest-frame, hard-band luminosity larger than 10^{43} erg/s. The luminosity range puts all these sources in the Seyfert regime, far from normal star forming galaxies. In principle, we may expect obscured star formation at a high level, up to 1000 $M_\odot$/year. Using the average $L_{2-10keV} - SFR$ relation in the redshift range

Table 5. X-ray properties of the 14 protocluster members

ID	N_H 10^{22} cm^{-2}	Γ	EW keV	$L_X(0.5-2keV)$ 10^{44} erg/s	$L_X(2-10keV)$ 10^{44} erg/s
7	$40.1^{+9.7}_{-8.4}$	$2.00^{+0.29}_{-0.28}$	0	3.47 ± 0.33	4.01 ± 0.27
36	$1.13^{+1.16}_{-1.05}$	$1.86^{+0.15}_{-0.14}$	$0.22^{+0.20}_{-0.17}$	1.27 ± 0.08	1.86 ± 0.15
57	< 1	$1.89^{+0.30}_{-0.28}$	0	0.21 ± 0.04	0.28 ± 0.07
58	$4.26^{+0.33}_{-0.32}$	$1.90^{+0.032}_{-0.031}$	$0.016^{+0.024}_{-0.016}$	31.92 ± 0.44	42.85 ± 0.68
74	< 2.7	$1.59^{+0.21}_{-0.20}$	$0.47^{+0.37}_{-0.31}$	0.17 ± 0.02	0.39 ± 0.07
75	$0.33^{+0.65}_{-0.32}$	$1.83^{+0.10}_{-0.09}$	$0.07^{+0.15}_{-0.07}$	2.05 ± 0.09	3.11 ± 0.17
80	$18.0^{+4.6}_{-3.9}$	$1.96^{+0.23}_{-0.20}$	$0.34^{+0.13}_{-0.12}$	3.26 ± 0.21	4.15 ± 0.25
86	< 2 (> 150)*	1.10 ± 0.20	$0.51^{+0.38}_{-0.32}$	0.07 ± 0.01 †	0.34 ± 0.05 †
9	< 2.2	1.80	–	0.086 ± 0.020	0.135 ± 0.051
12	$19.4^{+19.8}_{-19.4}$	1.80	–	0.12 ± 0.04	0.18 ± 0.10
34	0 (> 150)*	$-0.76^{+0.70}_{-0.84}$	–	< 0.01 †	0.026 ± 0.009 †
73	$13.3^{+7.6}_{-6.1}$	1.80	–	0.15 ± 0.05	0.23 ± 0.05
87	0	2.2 ± 0.4	–	0.18 ± 0.04	0.15 ± 0.09
90	$14.8^{+7.5}_{-5.8}$	1.80	–	0.18 ± 0.04	0.28 ± 0.07

Notes. X-ray properties from spectral fit for the 14 protocluster members identified in the X-ray, including the Spiderweb Galaxy (XID 58). The two sources flagged with the symbol * are considered Compton-thick candidates, and therefore have an estimated intrinsic absorption $> 1.5 \times 10^{24}$ cm^{-2}. The corresponding luminosity values, flagged with the symbol †, correspond to the observed luminosity, and not to the intrinsic, unabsorbed values, as for the other sources.

2.0-2.5 as measured from deep fields by Lehmer et al. (2016), we obtain $L_{2-10keV} \sim 10^{42}$ and $\sim 10^{43}$ erg/s for a SFR of 100 and 1000 $M_\odot$/yr, respectively. On the other hand, sources with high intrinsic absorption are unambiguously dominated by nuclear emission with absorption due to an optically thick torus, while absorption at the level of $N_H \sim 10^{22}$ cm^{-2} and below, can be associated to the interstellar medium of the host, if sufficiently dense (but see recents results in Norman et al. 2021, where column densities of $N_H \sim 10^{23}$ cm^{-2} can be reached at $z \sim 2$). In Figure 16 we plot the intrinsic N_H as a function of $L_{2-10keV}$ and show with green lines the corner where N_H and $L_{2-10keV}$ are consistent with being powered by star formation a the level of 100 and 1000 $M_\odot$/yr. We conclude that all the X-ray sources detected in the protocluster are consistent with being in the Seyfert and QSO regime.

We make an attempt to detect variations in the average properties of the X-ray protocluster members as a function of the distance from the Spiderweb galaxy. We find that half of the spectroscopically confirmed protocluster members is distributed within a radius of 90 arcsec (corresponding to $\sim$ 760 kpc). Therefore, we consider the properties of the X-ray emitting protocluster members below and above this radius. As already shown in Figure 9, the distribution of the X-ray sources is very similar to that of all the spectroscopically confirmed members. The fraction within and beyond 90 arcsec is $14 \pm 5\%$ and $13 \pm 5\%$, respectively. If we consider only protocluster members with $log(M_*) > 10.5$ (see Section 8) we find $33 \pm 13\%$ and $29 \pm 12\%$, within and beyond 90 arcsec, respectively. The X-ray properties do not change either. The distribution of intrinsic absorption in the two subsample is almost exactly symmetric (with one C-thick candidate per side) apart for source ID=7, that shows the highest measured N_H and is the closest to the Spiderweb. Also, the average soft (hard) band luminosity is $0.87\,(1.21) \times 10^{44}$ erg/s and $0.82\,(1.1) \times 10^{44}$ erg/s, within and beyond 90 arcsec, respectively. Our preliminary investigation shows, if any, marginally higher X-ray fraction, intrinsic absorption, and lower luminosities towards the protocluster center. However, this trend, if present, can be reliably measured only with a much larger statistics, implying a population studies of protoclusters with comparable X-ray depth.

Finally, we note that there are no protocluster members which have been identified as AGN from the optical and NIR spectrum beside the X-ray detected sources. A detailed discussion of the single sources can be found in Appendix C.

6.5. Average X-ray properties of X-ray silent, spectroscopic and color-selected protocluster members

We investigate the presence of X-ray emission below our detection threshold in the other spectroscopically identified protocluster members. We extract a list of 82 sources with spectroscopic redshift in the range $2.0 < z < 2.3$ after removing the sources already detected in the X-ray band, and all the sources embedded in the diffuse emission close to the Spiderweb galaxy within a radius of 12 arcsec (the so called "flies"). Then we use the background images already used to compute the sky coverage, where all the 107 identified X-ray sources have been previously removed. These soft and hard-band "background-only" images have been obtained simply by cutting out the X-ray detected sources and replacing the signal at their position with random noise consistent with the signal in the annular region surrounding it. These images, therefore, contain only the sky X-ray emission from sources below our detection threshold. Then, we simply stack the images at the position of the 82 optical sources previously selected. The resulting images, shown in Figure 17, provide an estimate of the cumulative emission from spectroscopically confirmed, non X-ray detected protocluster members in the soft and hard band. If we perform a simple aperture photometry in a circle of 2 arcsec (corresponding to a reasonable photometric aperture enclosing 95% of the flux in our field), we do not obtain significant detection in either band, formally measuring 39 ± 30 and 49 ± 40 in the soft and hard band, respectively. However, in the total (0.5 - 7 keV) band we have 89 ± 50 net counts, which formally has a statistical significance of $\sim 2\,\sigma$. If we consider this



Fig. 13. Top panel: the differential, normalized distribution of the soft fluxes in the Spiderweb Galaxy Field. The total sample of X-ray sources is shown with a thin line, while the subsample of the 14 protocluster members are shown with a thick line. Bottom panel: same as in the top panel in the hard band.



Fig. 14. The red histogram shows the distribution of the intrinsic absorption of the 8 protocluster members (including the Spiderweb) for which a robust measurement of N_H has been possible. The blue dashed histogram shows the upper limits on N_H obtained for 6 sources. The black solid line shows the distribution of N_H expected in the model of Gilli et al. (2007) at the flux limit of our observations. The two Compton-thick candidates, with a reflection-dominated spectrum, are not included.



Fig. 15. The distribution of the X-ray luminosity of the 11 protocluster members (excluding the Spiderweb and the two Compton-thick candidates) in the soft (thin red line), hard (thin blue line) and total band (thick magenta line). For clarity, the histograms for the soft and hard luminosities are shifted by 0.05 and 0.1 dex, respectively.

as a marginal detection, after correction for average vignetting ($\sim 5\%$), we obtain a soft-band flux of $\sim 6.2 \times 10^{-18}$ erg/s/cm^{-2} and a hard-band flux of $\sim 1.6 \times 10^{-17}$ erg/s/cm^{-2} per source. Assuming a typical spectrum with $\Gamma = 1.8$ and $N_{\rm H} \sim 10^{22}$ cm^{-2}, this gives intrinsic, rest-frame luminosity range of $2.3-3.0\times10^{41}$ erg/s and $3.6 - 4.8 \times 10^{41}$ erg/s (after correcting for intrinsic and Galactic absorption) in the 0.5-2.0 and 2-10 keV band, respectively. The average, total band X-ray luminosity of these sources therefore is well below $\sim 10^{42}$ erg/s, which, in turn, corresponds to a maximum SFR significantly less than 100 M$_\odot$/yr per source. This shows us that, with current *Chandra* capabilities, we can barely detect the average X-ray signal from spectroscopically confirmed members at the level of ~ 1 net count in our deep 715 ks exposure. At the same time, this would imply a widespread star formation activity at the level of few$\times$10 M$_\odot$/yr on average for all the protocluster members, while optical and NIR spectroscopy do not confirm this. Therefore, if such a weak X-ray signal could be confirmed, it may be ascribed to the combination of low-level nuclear activity and moderate star formation. Clearly, to explore the low X-ray luminosity regime of the bulk of the protocluster members, we should increase the current exposure by at least a factor of five. This would allow us to explore the star formation regime corresponding to $20-50$ M$_\odot$/yr in each single source, implying that all the strong-starburst galaxies may be detected. We conclude that X-ray silent spectroscopic members do not contribute to the AGN luminosity function of the protocluster, but they may be hosting starburst events or low-level AGN activity. This is currently out of *Chandra* capability (considering regular and large programs) and it definitely constitutes an interesting luminosity range to be explored by future, high-angular resolution X-ray missions, such as Lynx (The Lynx



Fig. 16. Scatter plot of N_H and $L_{2-10keV}$ for 11 sources (excluding the Spiderweb and the two Compton-thick candidates). Red circles correspond to measured N_H values, while green triangles to 1σ upper limits. The green box shows the parameter space still consistent with star formation at the level of 1000 $M_\odot$/yr or 100 $M_\odot$/yr (vertical dashed line).



Fig. 17. Left panel: stacked image of the 82 spectroscopically confirmed protocluster members in the soft band. The size of the cutout is 20×20 arcsec, while the red circle shows the 2 arcsec radius region used to perform aperture photometry on the stacked image. Right panel: same as in the left panel in the hard band.

Team 2018) and AXIS (Mushotzky & AXIS Team 2019; Marchesi et al. 2020).

Then, we applied the same procedure to the 81 color or narrow-band selected sources that have no X-ray detection, and we find similar results, possibly with a more convincing detection in the soft band, as shown in Figure 18. The usual aperture photometry within a circle of 2 arcsec provides 60±30 net counts in the soft band, therefore a formal detection at 2σ. The soft band photometry corresponds to a soft flux of $(9.4 \pm 5) \times 10^{-18}$ erg/s/cm^{-2}, and, assuming the usual mildly absorbed power law and $z = 2.156$, to a luminosity of $(3.6 \pm 1.8) \times 10^{41}$ and $(5.6 \pm 2.8) \times 10^{41}$ erg/s per source in the soft and hard band, respectively. This luminosity range is typical of starburst galaxies with SFR of the order of $30 - 50$ $M_\odot$/yr. In the hard band we obtain only an upper limit (formally the photometry gives 13 ± 39 net counts), which can be translated in a 2σ upper limits of 4.0×10^{41} and 6.0×10^{41} erg/s per source in the soft and hard band, respectively. This tells us that the average sources are softer than the adopted, AGN-like model. Despite the small signal obtained in the soft band, we further split the 81 color or



Fig. 18. Left panel: stacked image of the 81 color or narrow-band selected protocluster member candidates in the soft band. The red circle shows the 2 arcsec radius region used to perform aperture photometry on the stacked image. Right panel: same as in the left panel in the hard band.

narrow-band selected sources without X-ray detection, in four subsamples, according to their selection criteria: H_α emitter, Ly_α emitter, ERO (Kurk et al. 2004b), and photo-z (Dannerbauer et al. 2014; Shimakawa et al. 2018; Tadaki et al. 2019). We find 1 σ positive photometry in the soft band for Ly_α (13) and ERO (29) selected sources, and 1.5 σ positive photometry for photo-z (30). In the hard band we have both positive and negative photometry. Therefore, the 2σ signal in the soft band is confirmed, and it is contributed mostly by the 30 protocluster member candidates selected by photo-z. However, we have no control on the contamination of this list of protocluster member candidates. This aspect, coupled to the large statistical uncertainty associated to stacked emission measurements, hampers any conclusion on the X-ray emission from color-selected protocluster member candidates, except that we do not expect them to host significant nuclear activity.

Our general conclusion is that there is no significant X-ray emission that we have missed so far among all the confirmed or candidate protocluster members that have not been X-ray detected. However, the obtained marginal detections confirm that potentially we are able to detect star formation in the protocluster galaxies at the level of $\sim 30 - 50$ $M_\odot$/yr, by stacking their X-ray emission. Additional work must be done before we can draw any strong conclusion, in particular it is necessary to increase the number of spectroscopically confirmed members by more than a factor of two. In this perspective, future observation of this target with JWST and further spectroscopic follow-up may play a strong synergy with the available X-ray data and reveal further relevant properties of the galaxy population in the Spiderweb protocluster.

7. X-ray luminosity function of protocluster members

To compute a cumulative luminosity function in the protocluster region, we need a comoving volume representative of our sample. If we consider the comoving volume corresponding to a circle of 5 arcmin centered on the Spiderweb Galaxy, and the difference in the comoving radial distance between $z = 2.11$ and $z = 2.2$, we find a value of 2.584×10^4 Mpc^3. The comoving mass corresponding to this volume in the assumed cosmology is $M_{\rm com} = 1.05 \times 10^{15} M_\odot$, which is in the range of predicted virial mass at $z = 0$ for the Spiderweb protocluster. Actually, this is rather a lower limit to the mass included within a radius of

5 arcmin, since the central part of the protocluster is definitely overdense with respect to the ambient cosmic density.

Secondly, we need an estimate of the intrinsic luminosity of the Compton-thick candidates. As we mentioned, the luminosity values listed in Table 5 for XID 86 and XID 34, simply refer the observed luminosity, that, in the case of a reflection dominated spectrum, is only a fraction of the intrinsic power. We conservatively assume that the observed hard (2-10 keV) luminosity is $\sim$ 10% of the intrinsic one (corresponding to the upper envelope of the values observed for local Compton-thick AGN, see Marchesi et al. 2019; Zhao et al. 2021), and that the soft-band luminosity is 1.6 times the hard-band luminosity, roughly corresponding to a power law spectrum with $\Gamma = 1.8$. This very crude approximation allows us to obtain a first-order estimate of the intrinsic luminosity of the two Compton-thick candidates, and, therefore, to have the soft and hard-band luminosities for all the 13 protocluster members.

Finally, we need to include the sky coverage correction in the luminosity function. We collect the weight factor corresponding to the soft and hard-band fluxes of each sources, similarly to what we did to compute the number counts. Then, we consider the smallest weight for each source, computed according to Equation 2. Given the relatively high luminosities of the protocluster members, we only have a small ($\sim$ 1.15) correction for the two faintest sources (XID 12 and XID 34). For the sake of clarity, we note that their S/N in the corresponding detection bands is $\sim$ 3, which is far enough from the detection limit, and, therefore, corresponds to a very mild sky coverage correction.

In Figure 19 we show the cumulative rest-frame luminosity in the soft and hard band, after excluding the Spiderweb Galaxy. In both cases we show it with and without the two Compton-thick candidates, given the large uncertainties in their estimated luminosities. We also plot the cumulative luminosity function predicted at $z \sim 2.2$ in the model of Gilli et al. (2007). As expected, the excess measured in the Spiderweb Protocluster volume is consistent with the excess previously measured in the number counts. The excess is always about an order of magnitude at the lowest end and increases with luminosity, reaching a factor of 20 and beyond at $L_X \sim 10^{44}$ erg/s in both bands. This is also clear in Figure 20 where we show the ratio of the X-ray AGN luminosity function in the Spiderweb protocluster to the field luminosity function in the field from the model by Gilli et al. (2007). To compute the ratio we include the Compton-thick candidates. Removing them will result in a lower excess, decreased by a factor of $\sim$ 1.5, as shown in Figure 19. As we already discussed, this excess is due to the combination of the average galaxy overdensity in the protocluster region and the enhancement factor. In Section 9 we will provide an estimate of the enhancement factor alone.

8. AGN fraction in the Spiderweb Protocluster

The measurement of the AGN fraction as a function of the environment is an important observable that can give insights on the SMBH feeding mechanisms. Several studies in the X-ray band have been performed for relatively low redshift in massive systems. Typically, the AGN fraction among cluster members increases with the distance from the center, and it reaches values comparable with the field (of the order of $\sim$ 1%) when considering the full X-ray luminosity range (10^{41}-10^{44} erg/s), or significantly lower than the field when focusing at luminosities $> 10^{43}$ erg/s (see Puccetti et al. 2020, and discussion therein). Overall, the X-ray nuclear emission in cluster galaxies (with the noticeable exception of the central galaxy) is expected to be triggered by interactions (see Ehlert et al. 2015), or by ram pressure in the outskirts (Poggianti et al. 2017), while several other mechanism are contributing in the field. The situation is expected to be significantly different in protoclusters, where the lower density of galaxies, the different velocity distribution, and the much larger availability of cold gas, may contribute to produce an AGN fraction significantly larger than in the field. Here we focus on the AGN fraction, and try to compare the results we obtained in the Spiderweb Complex with the literature, focusing on the best studied protoclusters.

The first aspect to notice is that the presence of activity (star formation and/or nuclear) in galaxies has some dependence on the environment and on the cosmic epoch at the same time. The two dependences can not be easily decoupled, since the main characteristics of dense environments change with epoch as well. Locally, it is well established that star formation and nuclear activity, at least at the bright end, are suppressed in dense environments such as massive, virialized halos. At the same time, the AGN fraction seems to increase in cluster progenitors at redshift $z \geq 2$ (Martini et al. 2013; Alberts et al. 2016; Kalita et al. 2021), while several studies have found intense star formation in at least some cluster cores at $z \geq 1.5$ (Tran et al. 2010; Brodwin et al. 2013; Santos et al. 2014, 2015; Coogan et al. 2018), suggesting an inversion of the SF-density anticorrelation observed in local clusters. An important aspect is the large diversity among the cluster population in the range $1.5 < z < 2.0$, with some of them hosting passive, evolved galaxies, and some a rapidly evolving, active population.

This is consistent with an increase (possibly a co-evolution) of SF and AGN in massive halos in the range $1.5 < z < 2.0$ when approaching their formation epoch. At present, we do not have a comprehensive understanding on how the connection between AGN activity and large-scale galaxy environment proceeds with redshift. This connection may be particularly effective when entering the regime of protoclusters. The deep *Chandra* observation of the Spiderweb Complex, presented in this work, gives us the possibility to measure the X-ray AGN fraction in a protocluster environment, and the X-ray AGN enhancement factor with respect to the field at the same redshift. Clearly, the member galaxy selection is key in this case, since protocluster members are usually picked up from a selection of Lyα emitters, or from color-color selected subsamples, potentially introducing a bias associated to the galaxy type. Here, a convenient parameter is the stellar mass.

We recall that the fraction of X-ray AGN among Lyα emitters is usually low, both in the field and in protoclusters (see, e.g., Malhotra et al. 2003; Zheng et al. 2016). However, several works have claimed for enhanced nuclear activity in massive protoclusters. Apart early claims in the Spiderweb, based, at the time, on 2 spectroscopically confirmed X-ray detected AGN and 3 X-ray member candidates (Pentericci et al. 2002), high AGN fraction at the level of $15 - 20\%$ has been found in 2QZ (Lehmer et al. 2013) at $z \sim 3$, DRC at $z \sim 4$ (Vito et al. 2020), and at lower redshift in Cl0218.3-0510 at $z = 1.62$ (Krishnan et al. 2017). Recently, AGN identified through optical spectroscopy of X-ray data in the *Planck*-selected protocluster at $z \sim 2.16$ have been shown to amount to $(20 \pm 10)\%$ of all the identified protocluster members (Polletta et al. 2021). In all cases, the limited statistics make the uncertainty on the AGN fraction significantly larger than $30 - 50\%$. On the other hand, other protoclusters in the range $2 < z < 3$ appear to have significantly lower fraction, implying an enhancement factor consistent with unity (Lehmer et al. 2009; Digby-North et al. 2010; Macuga et al. 2019). The



Fig. 19. Left panel: the red, solid line with the red-shadowed area shows the cumulative rest-frame, soft-band luminosity function of the AGN in the Spiderweb protocluster, normalized to the comoving volume defined by $2.11 < z < 2.20$ and a circle of 5 arcmin centered on the Spiderweb Galaxy. The cyan solid line with the pink-shadowed area is the same without the luminosity correction for the two C-thick candidates. The black solid line is the XLF in the field from the model of Gilli et al. (2007). Shaded areas correspond to $1\,\sigma$ uncertainty. The Spiderweb Galaxy is not included in the luminosity function. Right panel: same as left panel in the rest-frame, 2-10 keV hard band.



Fig. 20. Ratio of the soft and hard band X-ray AGN luminosity functions in the Spiderweb protocluster to the field luminosity function from the model by Gilli et al. (2007).



Fig. 21. Distribution of M_* among the protocluster members in the redshift range $2.11 < z < 2.20$ (red histogram) and among the X-ray emitting members (blue histogram). The first bin ($9.5 < \log(M_*) < 10.0$) is actually an upper limit where we include all the sources that do not have K_S band counterpart in our image.

fraction of X-ray emitting AGN in protoclusters measured so far in the literature are shown in Table 6.

To measure the AGN fraction in the Spiderweb protocluster, we need to control the selection in stellar mass. As detailed in Section 3, we estimated stellar masses M_* for about 80% of the sample. Among the 96 protocluster members ($2.11 < z < 2.20$), we have a robust measurement of M_* for 62 of them, while 34 are not detected. We assume a nominal upper limit of $\log(M_*/\mathrm{M}_\odot) = 10$ (which is irrelevant in this work). We note that most of the sources without K_S counterpart are from the COALAS survey. In fact, from Table 3 of Jin et al. (2021), we notice that half of them are not detected in the K_S band, while 90% of them are detected in the I band, and therefore these sources should correspond to relatively low mass galaxies.

The distribution of M_* in the protocluster is shown in Figure 21. Note that we include all the sources that do not have K_S band counterpart in our image in the first bin $9.5 < \log(M_*) < 10.0$. This is clearly an oversimplification that, however, does not have any effect on our results. Ideally, we may compute an AGN fraction as a function of M_*, but given the low statistics, we simply consider all the sources with $\log(M_*/\mathrm{M}_\odot) > 10.5$. For completeness, we briefly discuss preliminary results on the distribution of protocluster sources in the $L_X - M_*$ plane in Section D. Finally, we recall that there are no protocluster members which have been identified as AGN from the optical and NIR spectrum beside the X-ray detected sources. Therefore, excluding the Spiderweb Galaxy, we have 13 X-ray AGN (11 excluding the source

Table 6. X-ray AGN fraction in a sample of protoclusters

Protocluster	z	X-ray AGN fraction	Ref
Cl0218.3-0510	1.62	17^{+6}_{-5}%	Krishnan et al. (2017)
Spiderweb	2.156	25.5 ± 4.5%	This work
PHz G237.01	2.16	20 ± 10%	Polletta et al. (2021)
2QZ	2.23	17^{+16}_{-9}%	Lehmer et al. (2013)
HS1700	2.3	$2.9^{+2.9}_{-1.6}$%	Digby-North et al. (2010)
USS 1558-003	2.53	$2.0^{+2.6}_{-1.3}$%	Macuga et al. (2019)
SSA22	3.09	$5.1^{+6.8}_{-3.3}$%	Lehmer et al. (2009)
DRC	4.002	15^{+20}_{-10}%	Vito et al. (2020)

with photometric counterpart and the one with double counterpart) over a total of 44 sources, for an X-ray AGN fraction with $L_{2-10keV} > 10^{43}$ erg/s of 29.5% (25%). To bracket this value, we consider maximum of ∼ 28 additional members, two with X-ray emission, as discussed in Section 6.2. Their inclusion would bring the AGN fraction to 21%. Considering this as a lower limit, we can estimate the AGN fraction in the Spiderweb protocluster as 25.5 ± 4.5%, where the error bar account also for the maximum systematic uncertainty. We stress that the uncertainty we quote here, based on the photometric study of Pannella et al. (in preparation), also includes the effect of the different area coverage between X-ray data and spectroscopic or color/emission line selection. A detailed treatment of the effects of the protocluster member selection within 5 arcmin would imply a significant additional effort and would only mildly reduce the uncertainty we have estimated.

As shown in Table 6, this value is somewhat higher but consistent with the measurement from Krishnan et al. (2017) at $z = 1.62$ and to the highest values found at $2 < z < 4$ by Lehmer et al. (2013, 2QZ), and by Vito et al. (2020, DRC)[12]. Three different works, instead, find values in the range 2-5%, within an uncertainty of a factor of 2. We plot these values in Figure 22, where the measurement in this work is marked with a blue dot. We present this plot only to compare the different values obtained in the literature so far, with no attempt to explore possible behaviour with redshift. The main reason is that these values have been obtained from data with different X-ray depth and different selection for the protocluster membership. The first aspect is not an issue, since all the medium-deep *Chandra* data used in these works reach luminosity levels of ∼ 10^{43} erg/s in both bands. On the other hand, we lack of an homogeneous and complete selection of the protocluster members in the optical. As of today, the only robust constraint on the evolution of the X-ray AGN fraction is a clear increase from the low local values at the level of 1% to > 10% in a few protoclusters at $z > 2$. Similar, if not larger uncertainties, are found for the enhancement factor, that is discussed in the next Section.



Fig. 22. The fraction of X-ray AGN over the total number of identified protocluster members from Table 6. The Spiderweb Protocluster is marked with a blue dot. The plot is meant to show the measurements of AGN fraction available in the literature, and, necessarily, includes measurements for different populations and different stellar mass ranges.

9. AGN enhancement fraction in the Spiderweb Protocluster

To compare with the field AGN fraction, we refer to the few works in the literature that provide an estimate of this key quantity. Lehmer et al. (2009) estimate an average fraction of $1.9^{+2.6}_{-1.3}$% and $0.7^{+1.6}_{-0.6}$% from $z \sim 2-3.4$ Lyman Break Galaxies and $z = 3.1$ Lyα Emitters in the CDF + SSA22 Field, respectively, for $\log(M_*/M_\odot) > 10.5$. These values have been obtained for $\log(L_{8-32keV}) > 43.5$, that, for a power law with $\Gamma = 1.8$, corresponds to $\log(L_{0.5-10keV}) > 43.66$. Only 7 sources satisfy this threshold, or 9 if we consider the Compton-thick candidates and assume that the hard-band intrinsic luminosity is ten times the observed one. Therefore, the fraction of AGN with $\log(L_{0.5-10keV}) > 43.66$ in our spectroscopic sample is 18.0 ± 2.0%. In this case the enhancement factor for $\log(L_{0.5-10keV}) > 43.66$ in the Spiderweb protocluster turns out to be $f_{\rm enh} = 9.5^{+16}_{-4.0}$ or $f_{\rm enh} = 26^{+100}_{-16}$, depending on the assumed field value.

Another estimate of the field X-ray AGN fraction is provided by Lehmer et al. (2013), using an independent sample of $z = 2.216 - 2.248$ HAEs from HiZELS in the wide-area ∼ 1.6 deg^2 COSMOS survey field (Sobral et al. 2013). Lehmer et al. (2013) found 10 *Chandra* sources matched to the 210 HiZELS HAEs within the C-COSMOS footprint, giving an AGN fraction in the low density environment of $3.0^{+4.0}_{-2.0}$%. The COSMOS flux limit of 1.3×10^{-15} erg/s/cm^2 in the 0.5–10 keV (see Marchesi et al. 2016) corresponds to $L_{0.5-10keV} \sim 4 \times 10^{43}$ erg/s. This limit is equivalent to that previously used for a comparison with LBG

[12] Note that Vito et al. (2020) considers only X-ray selection among sub-mm detected galaxies, revising for example the value found in SSA22 to 50^{+39}_{-24}%.

and HAE sources in the CDF + SSA22 Field. In this case the enhancement factor turns out to be $6.0^{+9.0}_{-3.0}$.

Previous estimates of the enhancement factor range from 2.1 ± 0.7 (Krishnan et al. 2017), to $3.5^{+3.8}_{-2.2}$ (Lehmer et al. 2013) and $6^{+10.3}_{-3.6}$ (Lehmer et al. 2009), in broad agreement with our results. As previously mentioned, however, there are works that measure substantially lower X-ray AGN fractions in protoclusters, concluding that there is no enhancement with respect to the field (Digby-North et al. 2010; Macuga et al. 2019). This difference may simply reflect the diversity among protoclusters and their different dynamical stage, as expected in such a rapidly evolving phase.

In all these cases, the uncertainties on the AGN fraction in the field clearly dominate the error budget. Nevertheless, we find that we always obtain an enhancement factor in the range 5-20 for X-ray AGN with $L_{0.5-10keV} >$ few $\times 10^{43}$ erg/s and $\log(M_*/\mathrm{M}_\odot) > 10.5$. On the other hand, it is well known that the enhancement factor drops below 1 in all the virialized halos at relatively low redshift ($z < 1$), except in the cluster outskirts (Koulouridis & Bartalucci 2019). Therefore, we confirm the existence of a strong evolution of the X-ray AGN enhancement factor in the central regions of clusters and protoclusters with redshift. However, we do expect to see also a strong dependence on the environment, and therefore a large diversity among the observed protoclusters, as probably suggested by the different conclusions on the enhancement factor at $z > 2$ presented in the literature. Only disentangling the complex pattern due to the combination of environmental effects and cosmic evolution we can reach a comprehensive picture of the galaxy population evolution in clusters.

10. Future perspectives

There are several promising approach to extend this work and gain further insights into the evolution of nuclear activity in protoclusters and connect it to the dearth of activity in virialized, mature clusters. Beside the difficult characterization of the protocluster candidates, that we already discussed in the Introduction, the first, obvious key aspect is the selection of the protocluster members, that largely differ among the few cases presented in the literature. Ideally, the protocluster membership should have a well-defined completeness as a function of stellar mass and distance from the center. Stellar mass should be used to compute AGN fraction above a given mass limit (typically, $M_* >$ few $\times 10^{10} M_\odot$). The information on the stellar mass of the active galaxies may also provide clues on the physical mechanisms that trigger the activity. Krishnan et al. (2017) explore the distribution of the stellar mass of the X-ray emitting AGN in the field and in protoclusters, and find no differences. Therefore, they suggest that triggering mechanisms are similar in both environments and that the higher X-ray AGN fraction in protoclusters is simply due to the same mechanism occurring more frequently because of the denser environment. However, recently Monson et al. (2021) speculated that the enhanced AGN fraction in the protocluster SSA22 at $z = 3.1$ is due to the larger masses (a factor of 2) of the LBG with respect to the field, rather than an enhanced merger activity. This issue is relevant also for the process quenching the rapidly star forming galaxies that are observed as quiescent galaxies in the protocluster, forming the nascent red sequence mentioned. On the same line, a census of the nuclear activity in the radio band is needed to complete the picture offered by the X-ray data and provide robust constraints on the feedback processes in the member galaxies.

Focusing on the Spiderweb protocluster, we foresee the following next steps. First, an extension of the protocluster membership based on NIR images and eventual spectroscopic follow-up, to achieve a well-defined and complete stellar-mass function. Then, a detailed investigation of the archival high-resolution HST data, that, coupled to the sub-mm data, may provide clues on the role of interactions and/or the presence of cold gas in triggering nuclear activity. Finally, we plan to exploit new radio data in the L band with JVLA and GMRT to characterize the protocluster population in this band.

This effort is aiming at making the Spiderweb an ideal science case for the study of protoclusters. In this respect, the role of X-ray data with high-angular resolution will be key for the extension of this approach to a large sample. Currently, *Chandra* is the only facility that can provide data with the required quality. The future of this science case heavily depends on the future of high-resolution X-ray astronomy, currently embodied in the Lynx (The Lynx Team 2018) and AXIS (Mushotzky & AXIS Team 2019) mission concepts.

11. Conclusions

We present the deep (715 ks) X-ray observation of the radio galaxy J1140-2629 (the Spiderweb Galaxy) and the surrounding large-scale structure with *Chandra*. The deep X-ray data allow us to map the nuclear activity in a dense environment down to the low-luminosity end of the Seyfert galaxy population, and test whether dense large-scale structure environments enhance X-ray nuclear activity, a phenomenon that has been investigated only recently at such a high redshift and only with shallower X-ray data. In addition, the sensitivity and exquisite angular resolution of *Chandra*, allows us to characterize the extended emission around the Spiderweb Galaxy and possibly identify thermal X-ray emission from the ICM, a key piece of information to constrain both the dynamical state of the protocluster and the feedback mechanism at work in the Spiderweb complex, and ultimately, to provide an overview of the baryon cycling in such an active environment at $z \sim 2$. In this first paper we focus on the nuclear activity of the galaxies in the Spiderweb protocluster, exploiting the combination of our deep X–ray data in the field with the large body of multi-wavelength and spectroscopic data available from the literature. The main results and the data products used in this work are made available on the project webpage[13]. Our results can be summarized as follows:

- We identified 107 X-ray sources within a radius of 5 arcmin centered on the Spiderweb Galaxy, down to flux limits of 1.3×10^{-16} erg/s/cm^2 and 3.9×10^{-16} erg/s/cm^2 and in the soft and hard band, respectively. We compute the cumulative number counts which are found to be consistent with results from X-ray deep fields in the literature.
- We identify 13 X-ray sources with spectroscopically confirmed counterparts belonging to the Spiderweb Complex (including the Spiderweb Galaxy itself) and 1 with a color-selected member candidate. Since these sources are found in the redshift range $2.11 < z < 2.20$, we classify them as protocluster members (as opposed to member of the Spiderweb Complex, which includes a larger redshift window $2.0 < z < 2.3$). The X-ray emitting protocluster members are distributed over an approximately rectangular region with with ~ 3.2 and 1.3 Mpc by side.

[13] http://www.arcetri.inaf.it/spiderweb/

- The X-ray spectral analysis finds 5 strongly absorbed AGN ($N_H > 10^{23}$ cm^{-2}), 2 sources with flat spectra, that we classify as Compton-thick candidates with reflection-dominated spectrum, and 7 sources with moderate or low absorption. After correcting for Galactic and intrinsic absorption, and with an educated guess for the intrinsic luminosity of the two Compton-thick candidates, the rest-frame 0.5-10 keV luminosities are found to be above 10^{43} erg/s for all the sources. By combining intrinsic absorption and intrinsic luminosities, we conclude that all the X-ray emitting protocluster members are powered by nuclear activity. We do not find hints of a change of the X-ray properties of the protocluster members with the distance from the Spiderweb Galaxy.
- We stack the images of all the X-ray silent, spectroscopically confirmed members, finding a marginal detection corresponding to average luminosities below $\sim 10^{41}$ erg/s, showing that there is no significant nuclear activity at about one order of magnitude below our detection threshold. A slightly stronger detection is obtained for the stacked emission of the color or narrow-band selected members, consistent with being powered by star formation at the level of $30 - 50$ M$_\odot$/year, but the unknown contamination fraction does not allow us to derive robust conclusions on their nature.
- The X-ray luminosity function of AGN in the volume associated to the Spiderweb protocluster is more than an order of magnitude larger than in the field at the same redshift in the range $10^{43} < L_X < 10^{44}$ erg/s in both bands, and significantly flatter, implying an increasing excess at the bright end.
- For $L_{2-10keV} > 10^{43}$ erg/s, we measure an X-ray AGN fraction of 25.5±4.5% among galaxies with $\log(M_*/M_\odot) > 10.5$, which is higher but still consistent with the values found at least in 4 protoclusters at similar redshift.
- The measured AGN fraction in the Spiderweb protocluster at $L_{0.5-10keV} > 4 \times 10^{43}$ erg/s, corresponds to an enhancement factor with respect to the field that ranges from 5 to 20 (with a preferred value $f_{enh} = 6.0^{+9.0}_{-3.0}$) with respect to the COSMOS field at comparable redshifts and stellar mass range), suggesting a significant environmental effect triggering nuclear activity in the Spiderweb protocluster.

Unfortunately, the decreasing sensitivity of *Chandra* due to the contamination layer on the ACIS detector, makes future studies of this kind very hard, and we can realistically foresee that only a handful of new targets can be studied with comparable detail in the next future. Nevertheless, the confirmation of an enhanced X-ray AGN activity in protoclusters is an important ingredient to plan studies of high-z large-scale structure with future X-ray facilities. By exploiting the available multiwavelength dataset on the Spiderweb field, we plan to further explore the properties of the X-ray protocluster members to investigate the main physical mechanism responsible for triggering the X-ray emission.

Acknowledgements. This work was carried out during the ongoing COVID-19 pandemic. The authors would like to acknowledge the health workers all over the world for their role in fighting in the frontline of this crisis. We thank the anonymous referee for detailed comments and positive criticism that helped improving the quality of the paper. We thank Stefano Marchesi for useful discussions. P.T. and R.G. acknowledge financial contribution from the agreement ASI-INAF n.2017-14-H.0. M.N. acknowledges INAF-1.05.01.86.20. S.B. acknowledges partial financial support from the Indark INFN Grant. A.S. is supported by the FARE-MIUR grant 'ClustersXEuclid' R165SBKTMA and INFN InDark grant. AS, MP, VS, LDM are supported by the ERC-StG 'ClustersXCosmo' grant agreement 716762. HD acknowledges financial support from the AEI-MCINN under grant with reference PID2019-105776GB-I00/DOI:10.13039/501100011033, and from the ACIISI, Consejería de Economía, Conocimiento y Empleo del Gobierno de Canarias and the European Regional Development Fund (ERDF) under grant with reference PROID2020010107. We thank Hans Moritz Günther for help with the use of the MARX software. We thank Malgorzata Sobolewska for assistance during the Chandra observations. This work is based in part on data collected at Subaru Telescope and obtained from the SMOKA, which is operated by the Astronomy Data Center, National Astronomical Observatory of Japan, (Baba et al. 2002). This research has made use of the services of the ESO Science Archive Facility.

References

Alberts, S., Pope, A., Brodwin, M., et al. 2016, ApJ, 825, 72
Anderson, C. S. & Carilli, C. L. 2022, ApJ, in press
Athreya, R. M., Kapahi, V. K., McCarthy, P. J., & van Breugel, W. 1998, A&A, 329, 809
Baba, H., Yasuda, N., Ichikawa, S.-I., et al. 2002, in Astronomical Society of the Pacific Conference Series, Vol. 281, Astronomical Data Analysis Software and Systems XI, ed. D. A. Bohlender, D. Durand, & T. H. Handley, 298
Beers, T. C., Flynn, K., & Gebhardt, K. 1990, AJ, 100, 32
Bertin, E. 2006, in Astronomical Society of the Pacific Conference Series, Vol. 351, Astronomical Data Analysis Software and Systems XV, ed. C. Gabriel, C. Arviset, D. Ponz, & S. Enrique, 112
Bertin, E., Mellier, Y., Radovich, M., et al. 2002, in Astronomical Society of the Pacific Conference Series, Vol. 281, Astronomical Data Analysis Software and Systems XI, ed. D. A. Bohlender, D. Durand, & T. H. Handley, 228
Branchesi, M., Gioia, I. M., Fanti, C., Fanti, R., & Cappelluti, N. 2007, A&A, 462, 449
Brodwin, M., Stanford, S. A., Gonzalez, A. H., et al. 2013, ApJ, 779, 138
Bruzual, G. & Charlot, S. 2003, MNRAS, 344, 1000
Calzetti, D., Armus, L., Bohlin, R. C., et al. 2000, ApJ, 533, 682
Carilli, C. L. & Anderson, C. S. 2022, ApJ, in press
Carilli, C. L., Röttgering, H. J. A., van Ojik, R., Miley, G. K., & van Breugel, W. J. M. 1997, ApJS, 109, 1
Champagne, J. B., Casey, C. M., Zavala, J. A., et al. 2021, ApJ, 913, 110
Coogan, R. T., Daddi, E., Sargent, M. T., et al. 2018, MNRAS, 479, 703
Croft, S., Kurk, J., van Breugel, W., et al. 2005, AJ, 130, 867
Dannerbauer, H., Kurk, J. D., De Breuck, C., et al. 2014, A&A, 570, A55
Digby-North, J. A., Nandra, K., Laird, E. S., et al. 2010, MNRAS, 407, 846
Doherty, M., Tanaka, M., De Breuck, C., et al. 2010, A&A, 509, A83
Ehlert, S., Allen, S. W., Brandt, W. N., et al. 2015, MNRAS, 446, 2709
Emonts, B. H. C., Lehnert, M. D., Dannerbauer, H., et al. 2018, MNRAS, 477, L60
Fan, X. 2006, New A Rev., 50, 665
Gabasch, A., Salvato, M., Saglia, R. P., et al. 2004, ApJ, 616, L83
Galametz, A., Stern, D., De Breuck, C., et al. 2012, ApJ, 749, 169
Giacconi, R., Rosati, P., Tozzi, P., et al. 2001, ApJ, 551, 624
Gilli, R., Comastri, A., & Hasinger, G. 2007, A&A, 463, 79
Gilli, R., Mignoli, M., Peca, A., et al. 2019, A&A, 632, A26
Gilli, R., Su, J., Norman, C., et al. 2011, ApJ, 730, L28
Gobat, R., Daddi, E., Onodera, M., et al. 2011, A&A, 526, A133
Hasinger, G., Miyaji, T., & Schmidt, M. 2005, A&A, 441, 417
Hatch, N. A., Overzier, R. A., Kurk, J. D., et al. 2009, MNRAS, 395, 114
HI4PI Collaboration, Ben Bekhti, N., Flöer, L., et al. 2016, A&A, 594, A116
Ilbert, O., Salvato, M., Le Floc'h, E., et al. 2010, ApJ, 709, 644
Iwasawa, K., Comastri, A., Vignali, C., et al. 2020, A&A, 639, A51
Jin, S., Dannerbauer, H., Emonts, B., et al. 2021, A&A, 652, A11
Kalita, B. S., Daddi, E., Coogan, R. T., et al. 2021, MNRAS, 503, 1174
Kocevski, D. D., Lubin, L. M., Gal, R., et al. 2009, ApJ, 690, 295
Kodama, T., Tanaka, I., Kajisawa, M., et al. 2007, MNRAS, 377, 1717
Komatsu, E., Smith, K. M., Dunkley, J., et al. 2011, ApJS, 192, 18
Koulouridis, E. & Bartalucci, I. 2019, A&A, 623, L10
Koyama, Y., Kodama, T., Tadaki, K.-i., et al. 2013, MNRAS, 428, 1551
Kriek, M., van Dokkum, P. G., Labbé, I., et al. 2009, ApJ, 700, 221
Krishnan, C., Hatch, N. A., Almaini, O., et al. 2017, MNRAS, 470, 2170
Kuiper, E., Hatch, N. A., Miley, G. K., et al. 2011, MNRAS, 415, 2245
Kurk, J. D., Pentericci, L., Overzier, R. A., Röttgering, H. J. A., & Miley, G. K. 2004a, A&A, 428, 817
Kurk, J. D., Pentericci, L., Röttgering, H. J. A., & Miley, G. K. 2004b, A&A, 428, 793
Kurk, J. D., Röttgering, H. J. A., Pentericci, L., et al. 2000, A&A, 358, L1
Lehmer, B. D., Alexander, D. M., Geach, J. E., et al. 2009, ApJ, 691, 687
Lehmer, B. D., Basu-Zych, A. R., Mineo, S., et al. 2016, ApJ, 825, 7
Lehmer, B. D., Lucy, A. B., Alexander, D. M., et al. 2013, ApJ, 765, 87
Liu, T., Tozzi, P., Wang, J.-X., et al. 2017, ApJS, 232, 8
Luo, B., Brandt, W. N., Xue, Y. Q., et al. 2017, ApJS, 228, 2
Macuga, M., Martini, P., Miller, E. D., et al. 2019, ApJ, 874, 54
Malhotra, S., Wang, J. X., Rhoads, J. E., Heckman, T. M., & Norman, C. A. 2003, ApJ, 585, L25
Marchesi, S., Ajello, M., Zhao, X., et al. 2019, ApJ, 872, 8

Marchesi, S., Civano, F., Elvis, M., et al. 2016, ApJ, 817, 34
Marchesi, S., Gilli, R., Lanzuisi, G., et al. 2020, A&A, 642, A184
Martini, P., Kelson, D. D., Kim, E., Mulchaey, J. S., & Athey, A. A. 2006, ApJ, 644, 116
Martini, P., Miller, E. D., Brodwin, M., et al. 2013, ApJ, 768, 1
Mayo, J. H., Vernet, J., De Breuck, C., et al. 2012, A&A, 539, A33
Miley, G. K., Overzier, R. A., Zirm, A. W., et al. 2006, ApJ, 650, L29
Monson, E. B., Lehmer, B. D., Doore, K., et al. 2021, ApJ, 919, 51
Muldrew, S. I., Hatch, N. A., & Cooke, E. A. 2015, MNRAS, 452, 2528
Mullaney, J. R., Daddi, E., Béthermin, M., et al. 2012a, ApJ, 753, L30
Mullaney, J. R., Pannella, M., Daddi, E., et al. 2012b, MNRAS, 419, 95
Mushotzky, R. & AXIS Team. 2019, in The Space Astrophysics Landscape for the 2020s and Beyond, Vol. 2135, 5025
Nesvadba, N. P. H., De Breuck, C., Lehnert, M. D., et al. 2011, A&A, 525, A43
Nonino, M., Dickinson, M., Rosati, P., et al. 2009, ApJS, 183, 244
Norman, C., Hasinger, G., Giacconi, R., et al. 2002, ApJ, 571, 218
Osterbrock, D. E. 1989, Annals of the New York Academy of Sciences, 571, 99
Overzier, R. A. 2016, A&A Rev., 24, 14
Pannella, M., Elbaz, D., Daddi, E., et al. 2015, ApJ, 807, 141
Pentericci, L., Kurk, J. D., Carilli, C. L., et al. 2002, A&A, 396, 109
Pentericci, L., Kurk, J. D., Röttgering, H. J. A., et al. 2000, A&A, 361, L25
Pentericci, L., Roettgering, H. J. A., Miley, G. K., Carilli, C. L., & McCarthy, P. 1997, A&A, 326, 580
Poggianti, B. M., Jaffé, Y. L., Moretti, A., et al. 2017, Nature, 548, 304
Polletta, M., Soucail, G., Dole, H., et al. 2021, A&A, 654, A121
Puccetti, S., Fiore, F., Bongiorno, A., et al. 2020, A&A, 634, A137
Puccetti, S., Vignali, C., Cappelluti, N., et al. 2009, ApJS, 185, 586
Richards, G. T., Strauss, M. A., Fan, X., et al. 2006, AJ, 131, 2766
Rigby, E. E., Hatch, N. A., Röttgering, H. J. A., et al. 2014, MNRAS, 437, 1882
Roettgering, H. J. A., Lacy, M., Miley, G. K., Chambers, K. C., & Saunders, R. 1994, A&AS, 108, 79
Roettgering, H. J. A., van Ojik, R., Miley, G. K., et al. 1997, A&A, 326, 505
Rosati, P., Tozzi, P., Giacconi, R., et al. 2002, ApJ, 566, 667
Salpeter, E. E. 1955, ApJ, 121, 161
Santos, J. S., Altieri, B., Tanaka, M., et al. 2014, MNRAS, 438, 2565
Santos, J. S., Altieri, B., Valtchanov, I., et al. 2015, MNRAS, 447, L65
Seymour, N., Stern, D., De Breuck, C., et al. 2007, ApJS, 171, 353
Shimakawa, R., Kodama, T., Tadaki, K. I., et al. 2014, MNRAS, 441, L1
Shimakawa, R., Koyama, Y., Röttgering, H. J. A., et al. 2018, MNRAS, 481, 5630
Sobral, D., Smail, I., Best, P. N., et al. 2013, MNRAS, 428, 1128
Tadaki, K.-i., Kodama, T., Hayashi, M., et al. 2019, PASJ, 71, 40
Tanaka, M., Toft, S., Marchesini, D., et al. 2013, ApJ, 772, 113
The Lynx Team. 2018, arXiv e-prints, arXiv:1809.09642
Tozzi, P., Gilli, R., Mainieri, V., et al. 2006, A&A, 451, 457
Tozzi, P., Rosati, P., Nonino, M., et al. 2001, ApJ, 562, 42
Tran, K.-V. H., Papovich, C., Saintonge, A., et al. 2010, ApJ, 719, L126
Valentino, F., Daddi, E., Finoguenov, A., et al. 2016, ApJ, 829, 53
Valtchanov, I., Altieri, B., Berta, S., et al. 2013, MNRAS, 436, 2505
van Ojik, R. 1995, PhD thesis, Leiden University, Netherlands
Vito, F., Brandt, W. N., Lehmer, B. D., et al. 2020, A&A, 642, A149
Wang, T., Elbaz, D., Daddi, E., et al. 2016, ApJ, 828, 56
Zhao, X., Marchesi, S., Ajello, M., et al. 2021, A&A, 650, A57
Zheng, Z.-Y., Malhotra, S., Rhoads, J. E., et al. 2016, ApJS, 226, 23
Zirm, A. W., Stanford, S. A., Postman, M., et al. 2008, ApJ, 680, 224
Zirm, A. W., Toft, S., & Tanaka, M. 2012, ApJ, 744, 181

Appendix A: Source list

In Table A we list the properties of the 107 X-ray sources detected in the FOV included in a circle of 5 arcmin centered on the Spiderweb Galaxy. Soft (0.5-2 keV) and hard (2-7 keV) net counts are obtained with aperture photometry. Values have been rounded to the first decimal digit except a few cases. Negative values are sometimes obtained due to background fluctuations. Soft and hard (2-10 keV) band energy fluxes are obtained multiplying the net counts, corrected for vignetting, by the corresponding conversion factors listed in Table 4. The errors are obtained combining independently the statistical photometric error with the systematic uncertainty on the spectral shape (hence, on the conversion factors). In case the S/N in a given band does not reach our threshold value $S/N > 2$, we quote the $2\,\sigma$ upper limit to the energy flux. The original `ASCII` file of Table 1 can be retrieved from the webpage[14].

Table A.1. Catalog of the X-ray detected sources within 5 arcmin from the Spiderweb Galaxy.

XID	RA	DEC	Cts/s (soft)	Cts/s (hard)	F_S 10^{-16} erg/s/cm^2	F_H 10^{-16} erg/s/cm^2
1	175.18593	-26.54934	26.2 ± 6.6	164.0 ± 13.8	4.3 ± 1.1	71.4 ± 11.2
2	175.27884	-26.50376	174.8 ± 13.9	246.6 ± 17.5	34.7 ± 3.8	121.2 ± 18.1
3	175.24638	-26.50113	6.9 ± 4.0	19.4 ± 6.1	< 2.0	7.6 ± 2.6
4	175.22588	-26.49542	220.5 ± 15.2	174.6 ± 13.7	29.3 ± 3.0	69.2 ± 10.6
5	175.20786	-26.49363	29.0 ± 5.9	28.1 ± 6.0	3.7 ± 0.8	11.0 ± 2.8
6	175.22015	-26.48645	241.1 ± 15.8	189.1 ± 14.2	31.9 ± 3.2	75.1 ± 11.4
7	175.19974	-26.48512	132.6 ± 12.6	235.3 ± 16.0	17.1 ± 2.1	92.4 ± 13.7
8	175.28694	-26.47671	19.0 ± 5.0	12.9 ± 5.2	4.4 ± 1.2	7.6 ± 3.2
9	175.23038	-26.47326	23.5 ± 5.4	12.2 ± 4.7	3.1 ± 0.7	4.8 ± 1.9
10	175.25893	-26.51378	18.5 ± 5.2	−1.7 ± 4.3	2.9 ± 0.8	< 3.0
11	175.29327	-26.48855	7.3 ± 4.7	18.6 ± 7.1	< 3.8	10.4 ± 4.2
12	175.22990	-26.47804	15.2 ± 4.9	9.8 ± 5.2	2.0 ± 0.7	3.8 ± 2.1
13^{r3}	175.28102	-26.44954	8.3 ± 3.5	−4.6 ± 2.2	1.9 ± 0.8	< 1.0
14	175.13669	-26.53742	22.0 ± 6.3	6.7 ± 6.1	4.9 ± 1.5	< 10.0
15	175.12944	-26.49295	16.6 ± 5.6	3.2 ± 5.5	2.5 ± 0.9	< 6.0
16	175.23132	-26.54720	22.4 ± 5.9	38.1 ± 8.2	3.9 ± 1.1	16.7 ± 4.2
17^{r1}	175.18042	-26.53844	3.7 ± 3.1	11.5 ± 4.8	< 1.5	4.9 ± 2.1
18	175.21500	-26.52062	5.3 ± 3.7	14.5 ± 5.3	< 1.7	5.7 ± 2.2
19	175.17738	-26.51848	−2.4 ± 2.1	13.9 ± 5.4	< 0.3	5.6 ± 2.3
20	175.11499	-26.48742	−3.7 ± 3.8	24.8 ± 7.7	< 0.8	15.2 ± 5.1
21	175.21033	-26.55919	44.8 ± 7.7	56.6 ± 9.1	10.2 ± 1.9	33.4 ± 7.0
22	175.20374	-26.55444	141.0 ± 12.5	259.1 ± 16.9	25.2 ± 2.9	118.5 ± 17.4
23	175.18608	-26.54500	378.1 ± 19.9	263.3 ± 17.1	62.6 ± 5.7	119.6 ± 17.6
24	175.25270	-26.53977	83.4 ± 9.8	86.2 ± 10.7	15.2 ± 2.1	39.6 ± 7.2
25	175.26349	-26.53928	5.0 ± 4.0	25.6 ± 7.6	< 3.0	15.1 ± 4.9
26^{r1}	175.16380	-26.53818	9.9 ± 4.6	1.8 ± 4.5	1.6 ± 0.8	< 4.5
27	175.24773	-26.53220	619.8 ± 25.4	574.2 ± 24.6	109.5 ± 9.3	265.3 ± 36.7
28	175.24940	-26.53080	29.9 ± 7.2	32.2 ± 7.5	5.3 ± 1.3	14.8 ± 3.9
29	175.24535	-26.53001	28.6 ± 6.6	14.9 ± 6.1	5.1 ± 1.2	7.1 ± 3.0
30	175.27881	-26.52859	339.7 ± 18.7	337.3 ± 19.1	91.4 ± 8.4	218.5 ± 31.3
31	175.18436	-26.52488	107.7 ± 11.0	277.6 ± 17.0	14.7 ± 1.8	112.4 ± 16.3
32	175.17839	-26.51327	74.8 ± 9.0	98.4 ± 10.5	10.0 ± 1.4	39.7 ± 6.7
33	175.24586	-26.50403	2.7 ± 2.7	7.7 ± 4.2	< 1.0	3.0 ± 1.5
34	175.24149	-26.49342	1.1 ± 2.9	16.2 ± 5.5	< 1.0	6.4 ± 2.3
35	175.19250	-26.48165	10.3 ± 3.9	5.6 ± 3.8	1.34 ± 0.52	< 5.0
36	175.16551	-26.47922	260.8 ± 16.6	175.2 ± 14.0	35.7 ± 3.5	72.9 ± 11.2
37	175.18231	-26.46532	164.2 ± 13.3	359.2 ± 19.5	22.0 ± 2.4	146.5 ± 20.9
38	175.26686	-26.46100	−1.5 ± 2.1	8.1 ± 4.3	< 0.5	3.5 ± 1.7
39	175.26224	-26.45923	77.5 ± 9.4	19.0 ± 6.3	12.5 ± 1.8	8.3 ± 2.9
40	175.28162	-26.45306	16.0 ± 5.7	62.8 ± 9.4	3.6 ± 1.3	36.7 ± 7.3

[14] `http://www.arcetri.inaf.it/spiderweb/`

41	175.27260	-26.44206	13.2 ± 5.1	18.3 ± 6.8	3.0 ± 1.2	11.2 ± 4.4
42	175.16692	-26.43220	90.8 ± 10.8	101.6 ± 12.3	13.8 ± 1.9	42.5 ± 7.6
43	175.20650	-26.42832	119.2 ± 12.0	28.0 ± 7.7	17.6 ± 2.2	11.5 ± 3.5
44	175.14221	-26.54348	1.6 ± 4.7	24.5 ± 7.7	< 2.4	14.4 ± 4.9
45	175.15338	-26.53828	8.9 ± 5.3	14.9 ± 6.7	< 3.2	6.4 ± 3.0
46	175.13422	-26.51784	7.7 ± 5.0	12.6 ± 6.7	< 2.8	5.5 ± 2.7
47	175.26782	-26.45793	3.6 ± 4.4	11.6 ± 6.4	< 2.0	4.9 ± 2.5
48	175.15282	-26.45285	12.4 ± 5.1	22.5 ± 6.8	1.86 ± 0.77	9.7 ± 3.2
49	175.16095	-26.44319	7.5 ± 4.4	11.0 ± 5.5	< 2.3	4.6 ± 2.4
50	175.14491	-26.44095	2.2 ± 5.3	51.5 ± 9.7	< 2.0	22.2 ± 5.1
51	175.23083	-26.43491	11.7 ± 4.9	8.7 ± 5.9	1.68 ± 0.70	3.5 ± 2.3
52	175.20361	-26.43283	6.3 ± 4.3	68.0 ± 9.3	< 2.0	27.6 ± 5.3
53	175.19453	-26.42296	1.2 ± 4.0	44.7 ± 8.6	< 1.4	18.3 ± 4.3
54	175.13251	-26.46288	3.3 ± 4.8	17.4 ± 7.3	1.9	7.2 ± 3.2
55	175.18709	-26.50135	1.4 ± 2.7	38.6 ± 6.8	< 0.8	16.0 ± 3.5
56	175.18121	-26.47711	4.9 ± 3.0	34.1 ± 6.6	< 1.5	14.5 ± 4.0
57	175.15547	-26.50483	41.2 ± 7.1	26.6 ± 6.4	5.6 ± 1.1	10.8 ± 3.0
58	175.20148	-26.48577	5355.1 ± 73.4	4033.0 ± 63.7	689.6 ± 52.0	1575.0 ± 208.8
59	175.27609	-26.48332	4.0 ± 4.3	20.6 ± 6.9	< 2.0	8.9 ± 3.2
60	175.22218	-26.43210	9.9 ± 4.6	17.5 ± 6.4	1.43 ± 0.68	7.1 ± 2.7
61	175.20068	-26.43743	8.2 ± 4.1	23.8 ± 6.7	1.15 ± 0.58	9.7 ± 3.0
62	175.19254	-26.43634	7.9 ± 4.0	7.9 ± 4.6	1.12 ± 0.55	3.2 ± 1.9
63	175.22252	-26.45065	26.4 ± 5.9	49.4 ± 8.0	3.6 ± 0.8	19.7 ± 4.1
64	175.25787	-26.44586	30.0 ± 6.8	14.0 ± 6.7	4.7 ± 1.1	5.8 ± 2.9
65	175.27296	-26.44957	21.9 ± 6.0	22.6 ± 7.2	4.2 ± 1.2	11.2 ± 3.9
66	175.27232	-26.45238	9.7 ± 4.5	23.9 ± 7.8	1.73 ± 0.81	11.1 ± 3.9
67	175.13544	-26.52808	53.3 ± 8.3	53.8 ± 9.2	10.2 ± 1.8	27.9 ± 6.0
68	175.14703	-26.47453	3.7 ± 3.9	18.9 ± 6.4	< 1.6	7.7 ± 2.8
69	175.16040	-26.46140	5.3 ± 3.9	29.6 ± 6.8	< 1.8	12.0 ± 3.2
70	175.16412	-26.46660	11.2 ± 4.3	13.3 ± 5.0	1.50 ± 0.59	5.4 ± 2.1
71	175.16164	-26.48617	73.7 ± 9.0	17.4 ± 5.3	9.9 ± 1.4	7.1 ± 2.4
72	175.18102	-26.46208	20.4 ± 5.1	11.2 ± 4.6	2.8 ± 0.7	4.6 ± 2.0
73	175.18962	-26.46955	12.5 ± 4.2	24.9 ± 5.7	1.68 ± 0.57	10.3 ± 2.7
74	175.18440	-26.48541	60.7 ± 8.0	39.7 ± 6.8	8.5 ± 1.3	17.0 ± 3.7
75	175.19153	-26.48805	495.0 ± 22.4	334.8 ± 18.7	64.2 ± 5.6	133.0 ± 19.0
76	175.20314	-26.46463	1.1 ± 2.5	11.8 ± 4.5	< 0.8	4.8 ± 1.9
77	175.22797	-26.45959	5.0 ± 3.4	42.3 ± 7.4	< 1.6	17.2 ± 3.8
78	175.23540	-26.46467	14.4 ± 4.6	9.8 ± 4.7	1.90 ± 0.63	3.8 ± 1.9
79	175.25610	-26.46475	0.5 ± 3.2	24.4 ± 6.6	< 1.1	10.9 ± 3.3
80	175.25993	-26.46278	248.4 ± 16.1	314.3 ± 18.6	41.2 ± 4.1	144.2 ± 20.8
81	175.20267	-26.48068	58.9 ± 8.0	43.6 ± 7.2	7.6 ± 1.2	17.1 ± 3.6
82	175.27940	-26.47438	10.1 ± 4.8	29.9 ± 7.7	1.7 ± 0.8	13.4 ± 3.9
83	175.14645	-26.49300	4.4 ± 3.6	38.8 ± 7.5	< 1.6	16.0 ± 3.7
84	175.15465	-26.49537	7.1 ± 4.0	15.2 ± 5.4	< 2.0	6.2 ± 2.4
85	175.16492	-26.49530	56.2 ± 8.0	100.1 ± 10.5	7.5 ± 1.2	40.7 ± 6.8
86	175.18535	-26.48916	51.6 ± 7.5	52.0 ± 7.8	7.1 ± 1.2	21.8 ± 4.4
87	175.19443	-26.48622	24.2 ± 5.4	6.3 ± 3.8	3.1 ± 0.7	2.5 ± 1.5
88	175.20360	-26.48595	37.7 ± 8.0	23.5 ± 6.2	4.9 ± 1.1	9.2 ± 2.7
89	175.21483	-26.48368	7.6 ± 3.5	42.7 ± 7.2	1.00 ± 0.46	16.9 ± 3.6
90	175.21136	-26.49230	25.2 ± 5.5	23.3 ± 5.6	3.25 ± 0.75	9.0 ± 2.5
91	175.22507	-26.49026	17.8 ± 4.8	20.5 ± 5.4	2.4 ± 0.7	8.1 ± 2.4
92	175.22772	-26.49112	89.5 ± 9.7	14.6 ± 4.9	11.8 ± 1.5	5.7 ± 2.1
93	175.23019	-26.49082	66.7 ± 8.5	34.6 ± 6.7	8.7 ± 1.3	13.5 ± 3.1
94[r1]	175.23413	-26.48683	5.2 ± 2.9	8.7 ± 3.8	< 1.3	3.4 ± 1.5
95	175.24232	-26.50758	47.2 ± 7.5	54.0 ± 8.5	6.3 ± 1.1	21.1 ± 4.3
96	175.26640	-26.51311	116.2 ± 11.3	344.1 ± 19.3	18.4 ± 2.3	146.2 ± 20.9
97	175.26569	-26.51449	13.4 ± 4.4	41.4 ± 7.4	2.1 ± 0.7	17.4 ± 3.9
98	175.24670	-26.53411	77.1 ± 10.0	99.3 ± 11.4	12.9 ± 1.9	43.1 ± 7.5
99	175.24873	-26.53539	53.5 ± 8.5	52.1 ± 8.9	8.8 ± 1.5	21.8 ± 4.7
100	175.22601	-26.56204	40.9 ± 7.2	36.2 ± 7.8	11.3 ± 2.2	25.8 ± 6.5
101	175.18367	-26.51827	14.0 ± 4.7	3.5 ± 3.9	1.9 ± 0.6	< 4.0
102	175.14857	-26.54114	163.1 ± 13.8	135.8 ± 13.8	31.4 ± 3.5	70.2 ± 11.7
103	175.16844	-26.55954	6.7 ± 5.1	61.5 ± 9.8	< 3.6	35.3 ± 7.3
104[r3]	175.20998	-26.56691	−1.3 ± 1.8	9.0 ± 4.2	< 0.7	6.8 ± 3.3

105	175.13370	-26.49614	-2.1 ± 1.5	7.5 ± 3.6	< 0.2	3.2 ± 1.6
106^{r3}	175.26247	-26.43484	8.3 ± 3.5	3.7 ± 3.6	1.61 ± 0.70	< 5.5
107	175.26004	-26.49588	2.0 ± 3.5	13.8 ± 5.6	< 1.3	5.9 ± 2.5

Notes. The RA and DEC positions refer to the centroid of the X-ray detected emission. If a source does not satisfy $S/N > 2$ in one band, we report the $2\,\sigma$ upper limit. Sources with apex ^{r1} and ^{r3} have an extraction radius 1 and 3 pixels, respectively, smaller than r_{ext} as defined in Section 5.1.

Appendix B: Spectra of X-ray members

In Figure B.1, we show the folded spectra of the 8 X-ray members of the Spiderweb Protocluster that have at least 40 net counts in the soft (0.5-2 keV) or hard (2-7 keV) band, along with the best-fit spectral model. Spectra have been rebinned with a minimum of 20 counts per energy bin for display purpose. The spectrum of the Spiderweb Galaxy (XID 58) will be discussed in much greater detail in a companion paper (Tozzi et. al. in preparation). The spectral model used in all the fits consists in a fixed modelization of the Galactic absorption with the *Xspec* model `tbabs`, with $N_{\rm H\,gal} = 3.18 \times 10^{20}$ cm^{-2}, an intrinsic power law with a free slope Γ, an intrinsic absorption modeled with `zwabs`, and a Gaussian line component that is allowed to vary in a range $1.9 - 2.1$ keV, approximately centered on the energy 2.03 keV corresponding to the neutral iron K_α line at a rest frame energy of 6.4 keV.

For completeness, in Figure B.2, we show the folded spectra of the 6 X-ray members of the Spiderweb Protocluster that do not reach 40 in the soft and in the hard band. Spectra have been rebinned with a minimum of only 10 counts per energy bin for display purpose. The spectral model used in the fits is significantly simpler than the previous one, and consists in a fixed modelization of the Galactic absorption with the *Xspec* model `tbabs`, with $N_{\rm H\,gal} = 3.18 \times 10^{20}$ cm^{-2}, an intrinsic power law with a slope frozen to $\Gamma = 1.8$, and an intrinsic absorption modeled with `zwabs`. Note that for sources XID 34 and XID 87, the spectral slope is left free instead of the intrinsic absorption, because the simple Compton-thin model is not able to provide acceptable fits. Best-fit parameters for all the sources are shown in Table 5.



Fig. B.1. Folded spectra of the X-ray members of the Spiderweb protocluster with more than 40 net counts in the soft or in the hard band, including the Spiderweb Galaxy itself (XID 58).



Fig. B.1. - continued .



Fig. B.2. Folded spectra of the X-ray members of the Spiderweb protocluster with less than 40 net counts in the soft and in the hard band.

Appendix C: Notes on single sources

In this Appendix we briefly discuss specific features of all the X-ray detected members of the Spiderweb protocluster, except the Spiderweb Galaxy (XID 58).

Appendix C.1: XID 7

This source has been identified as a candidate Hα emitter in Kurk et al. (2004b) (ID=272), and a photometric redshift of $z = 2.18^{+0.08}_{-0.05}$ (ID=493) has been measured by Tanaka et al. (2013). Kuiper et al. (2011) investigated the NIR spectrum with SINFONI (see their ID 12), and found a spectroscopic redshift of $z = 2.1447 \pm 0.0023$ from the O[III] line, observed with a flux of $8.2 \pm 1.5 \times 10^{-17}$ erg/s/cm^2/Å. In the X-ray, this source shows a Compton thin spectrum with the highest intrinsic absorption in the sample ($\sim 4 \times 10^{23}$ cm^{-2}), and a luminosity in the QSO range ($L_{0.5-10keV} \sim 7.5 \times 10^{44}$ erg/s). It is the X-ray source closest to the Spiderweb Galaxy.

Appendix C.2: XID 9

The optical counterpart of source XID 9 has been originally selected as a candidate Lyα emitter. The optical spectrum is shown in Pentericci et al. (2002) (ID=1612), where the Lyα line has been measured to have a flux of 3.9×10^{-17} erg/s/cm^2, with an $EW > 30$ Å and a $FWHM = 490$ km/s This source has been confirmed also with Keck spectroscopy by Croft et al. (2005) (ID=L675) and the Lyα has been revised to have $FWHM = 200 \pm 30$ km/s with a flux of 6.4×10^{-17} erg/s/cm^2. The NV line is also detected with $FWHM = 200 \pm 30$ km/s for a flux of 4.1×10^{-17} erg/s/cm^2.

Appendix C.3: XID 12

The redshift of XID 12 (z=2.1628) has been obtained with Subaru in Shimakawa et al. (2014) (ID=71). The X-ray spectrum is consistent with negligible intrinsic absorption, while the total luminosity $L_{0.5-10keV} \sim 2.2 \times 10^{43}$ erg/s is the lowest among the X-ray protocluster member, but still in the Seyfert-like regime. This source has the largest offset from its optical counterpart, due to the extended nature of its emission as noted in the text. Possible association of the extended emission with radio jets will be investigated when the wide-field, high-resolution radio images will be available.

Appendix C.4: XID 34

This source has been identified with APEX LABOCA in Dannerbauer et al. (2014) (ID=DKB12a) with a flux $S_{870\mu} = 5.0 \pm 1.4$ mJy for a $S/N = 3.6$. It also has a strong detection at 24μ with $S_{24\mu} = 303.4 \pm 5.0$ μJy and significant detection in the range 160 - 500 μJy, but it only have an upper limit at 1.4 GHz with JVLA. There are 4 Hα emitters consistent with the LABOCA position, however DKB12a is the closest to the X-ray position. The Hα and FIR-SED derived SFR are 240 and 860 $M_\odot$/yr, respectively, suggesting the presence of dust. The total FIR luminosity is 5.0×10^{12} $L_\odot$. Eventually, a redshift of $z = 2.1690$ has been measured with Subaru in Shimakawa et al. (2014) (ID 29). Shimakawa et al. (2018) identified a possible subgroup at the position of this source. The source has been detected also in the CO line in the COALAS survey (Jin et al. 2021, see their Table 3), with an offset of 2 arcsec between the ATCA and X-ray centroids. The high FIR luminosity is consistent with the X-ray tentative classification as a Compton-thick AGN, due to an extremely flat (if not inverted) power law spectrum. Despite the very low flux, the estimated intrinsic luminosity would be $L_{2-10keV} \sim 2.6 \times 10^{43}$ erg/s in our simplistic assumption that the observed (reflected) luminosity is $\sim 10\%$ of the intrinsic emission.

Appendix C.5: XID 36

The optical counterpart of source XID 36 has been originally selected as a Lyα emitter with $B = 24.8$. The optical spectrum is shown in Pentericci et al. (2002), where the source has been classified as a QSO (ID=1687). Its spectrum shows Lyα, SiIV and CIV emission lines. The Lyα line has been measured to have a flux of 56×10^{-17} erg/s/cm^2, with an $EW = 164$ Å and a $FWHM = 5800$ km/s. The Lyα and CIV lines have a double–peaked profile due to absorption by neutral, metal-enriched gas. Presence of dust is suggested by the low Lyα/CIV line ratio compared to normal values for QSOs (Lyα/CIV $\sim$ 2.5 from composite spectra of QSO BLRs Osterbrock 1989).

The spectrum (ID=X3/L968d) obtained by Croft et al. (2005) found only moderate absorption blueward of Lyα, and corrected the redshift from $z = 2.183$ Pentericci et al. (2002) to $z = 2.162$. This AGN shows broad and narrow Lyα, broad and narrow CIV λ1549, and broad NV. The CIV emission was fitted with two Gaussian components, and the broad and narrow Lyα and NV were fitted with three Gaussian components. For the Lyα they found $FWHM = 4790 \pm 40$ km/s and $FWHM = 122 \pm 20$ km/s for the broad and narrow component, respectively. For the CIV they found $FWHM = 12180 \pm 280$ km/s and $FWHM = 1070 \pm 50$ km/s for the broad and narrow component, respectively. The NV line has a measured $FWHM = 4041 \pm 40$ km/s. The QSO nature of this source is confirmed by the high X-ray luminosity ($L_{0.5-10keV} \sim 3.1 \times 10^{44}$ erg/s). The source shows a moderate absorption in the X-ray at the level of $\sim 10^{22}$ cm^{-2}, but also shows hints of a strong Fe line with rest-frame EW $\sim$ 0.22 keV, despite with large uncertainties.

Appendix C.6: XID 57

The redshift of XID 57 (z=2.1426) has been obtained with Subaru in Shimakawa et al. (2014) (ID=14). The X-ray spectrum is typical of a Type I AGN with typical Seyfert luminosity $L_{0.5-10keV} \sim 5.0 \times 10^{43}$ erg/s.

Appendix C.7: XID 73

The source has been identified with an ERO candidate in Kurk et al. (2004b) with ID=463. It is the only X-ray emitting protocluster member in our catalog that has not been spectroscopically confirmed. In the X-ray appears as an obscured Seyfert galaxy.

Appendix C.8: XID 74

This source is classified as a quiescent galaxy despite it is detected in Hα probably due to the presence of the AGN that, however, does not contribute significantly to the overall SED. This source is investigated in detail in Doherty et al. (2010) where they measure it to have $z = 2.172$ (ID 456) and to be consistent with being formed in an intense burst of star formation, with a derived age of $1.6^{+1.1}_{-0.7}$ Gyr. The low current star formation rate $< 1.0\ M_{\odot}$/yr derived from the SED fit confirms that the galaxy is now in a quiescent phase. The X-ray spectrum is consistent with a Type I or moderately obscured Seyfert galaxy. Also, the neutral Fe emission line is detected with $EW \sim 0.5$ keV, despite with significant uncertainties.

Appendix C.9: XID 75

This source was selected as an Hα emitter candidate in Kurk et al. (2004b) and eventually spectroscopically confirmed (Kurk et al. 2004a) (ID=215). Its spectrum shows continuum emission plus a very broad line which almost covers the full spectral range, and it is identified with Hα. For this AGN the N[II] line is blended with the very broad Hα line and impossible to discern. The measured FWHM of the line is 5300 ± 800 km/s for a flux of $46.2 \pm 8.8 \times 10^{-17}$ erg/s/cm^2 and an $EW = 150 \pm 30$Å. The Hα emission is not considered therefore associated to star formation rate. The X-ray spectrum is typical of an unobscured QSO, with $L_{0.5-10keV} \sim 5.2 \times 10^{44}$ erg/s, consistent with the presence of the broad optical line.

Appendix C.10: XID 80

Source XID 80 has been identified with source ID 16 in Pentericci et al. (2002) as a candidate Lyα emitter encompassed by a very strong and extended Lyα emission, and therefore classified as an AGN. Eventually its spectrum has been obtained by Croft et al. (2005) at $z = 2.149$ (ID X16/L778d). The spectrum shows a broad Lyα (rest-frame deconvolved FWHM 890 km/s), and several emission lines such as NV (λ1240), CIV (λ1549), HeII (λ1640), CIII] (λ1909),[CII] (λ2326), and MgII (λ2798), all with $FWHM$ ranging from 1200 to 2000 km/s (see Table 2 of Croft et al. 2005).

This source has been also identified with APEX LABOCA at 870 μm with $S_{870\mu m} = 4.2 \pm 1.4$ mJy for a $S/N = 2.9$ in Dannerbauer et al. (2014) (ID=DKB16). Despite the faint signal, this source is clearly detected in IR and radio bands (from $S_{24\mu} = 572.1 \pm 5.0\ \mu$Jy to $S_{1.4GHz} = 76.2 \pm 19.0\ \mu$Jy with the VLA (see their Table 2). The estimated SFR is $\sim 1140 M_{\odot}$/yr and $\sim 830 M_{\odot}$/yr from the Hα and FIR SED, respectively, and the total FIR luminosity is measured to be $4.8 \times 10^{12}\ L_{\odot}$. The X-ray spectrum shows that this source is an obscured, Type II QSO, with with an unobscured luminosity $L_{0.5-10keV} \sim 7.4 \times 10^{44}$ erg/s.

Finally, this source has the second largest offset from its optical counterpart, due to a possible extended nature of the emission as noted in the text. Possible association of the extended emission with radio jets will be investigated when the wide-field, high-resolution radio images will be available.

Appendix C.11: XID 86

Source XID 86 is identified in Pentericci et al. (2002) (Source ID=5) with a faint extended ERO with $R - K = 5.8$ ($I - K = 5.3$) consistent with having a redshift close to the Spiderweb Galaxy. Eventually, the source has been spectroscopically confirmed by Croft et al. (2005) at $z = 2.162$ (ID 5Xd) Faint but significant Lyα emission (rest-frame deconvolved $FWHM \sim 400$ km/s) has been detected. It was not possible to classify this source as a starburst or an AGN, but the X-ray luminosity provides strong evidence for the presence of an AGN, as already noticed in the literature (Pentericci et al. 2002; Croft et al. 2005). In the X-ray, this source 86 shows a flat spectrum ($\Gamma \sim 1$), with hardness ratio $HR \sim 0$. We consider it a candidate Compton-thick source, with the X-ray spectrum dominated by a reflection component, and an estimated intrinsic luminosity $L_{2-10keV} \sim 3.4 \times 10^{44}$ erg/s. The detection of the neutral Fe emission line with $EW \sim 0.50$ keV strengthen the Compton-thick hypothesis.

Appendix C.12: XID 87

Source XID 87 has been selected as an ERO candidate in Kurk et al. (2004b) with ID=284 and identified as an Hα emitter in Shimakawa et al. (2018) (ID=48). The redshift has been measured thanks to a ALMA CO(3–2) observation presented in Tadaki et al. (2019). The CO(3-2) line has a redshift of 2.157 with a $FWHM = 232$ km/s. This source has been detected also in the CO(1-0) line in the COALAS survey (Jin et al. 2021, see their Table 3). The X-ray spectrum shows the steepest slope in the sample, and it is typical of a Type I AGN, with $L_{0.5-10keV} \sim 3.3 \times 10^{43}$ erg/s.

Appendix C.13: XID 90

This source was originally selected as an Hα emitter candidate in Kurk et al. (2004b) with ID=154. It has been detected also in the CO line in the COALAS survey (Jin et al. 2021, see their Table 3). We remark that the X-ray centroid is closer to a nearby source that, nevertheless, has similar color and preliminary photo-z consistent with being in the protocluster. This is the only X-ray source in our sample that has two possible counterparts, both consistent with being in the protocluster, but with only one spectroscopically confirmed. The X-ray spectrum is typical of Type II AGN, with a Seyfert-like unabsorbed luminosity of $L_{0.5-10keV} \sim 4.6 \times 10^{43}$ erg/s.

Appendix D: L_X-M_* relation

In Figure D.1 we show the $L_X - M_*$ plane for all the protocluster members with spectroscopic redshift. We plot 11 sources with X-ray emission, excluding the spiderweb and excluding also the two Compton-thick candidates, whose intrinsic luminosity cannot be estimated accurately. In addition, we show also the average X-ray luminosity per source that we obtain by stacking the sources without X-ray detection in 4 stellar mass bins. The mass bins have been chosen in order to have a similar number of sources (≥ 10) in each of them, except for the first bin ($9.5 < \log(M_*) < 10.0$) where we included all the sources without K_S-band counterpart. Before stacking the X-ray images at the source position, we removed the sources closer than 12 arcsec to the Spiderweb Galaxy, to avoid spurious signal due to the irregular diffuse emission. We have 38 sources in the first bin, then 13 sources in the $10.0 < \log(M_*) < 10.5$ bin, and 11 and 12 sources in the $10.5 < log(M_*) < 10.9$ and $10.9 < \log(M_*) < 11.6$ bin, respectively. Despite in all the cases we formally have a positive photometry, most of them are very close to zero, and, in all the cases, consistent with zero whitin 1 or 1.5 σ. Therefore, we must consider the blue points and the corresponding error bars as upper limits. The error bars on $\log(M_*)$ are ~ 0.2 for the X-ray detected sources, while corresponds to the bin width for the stacked points.

Apart from a very scattered, global trend of having higher intrinsic L_X at higher M_*, we note that the sources not detected in X-ray do not have luminosity close to our detection limit, but at least an order of magnitude lower, implying that they are not hosting AGN comparable to those hosted by the X-ray detected protocluster members. In the right panel of Figure D.1 we also plot, as a reference, the average L_X-M_* relation obtained in the field at $z \sim 2$ in Mullaney et al. (2012a) for the rest-frame 2-10 keV band. Apart from the expected scatter at high masses (see Mullaney et al. 2012b), that is entirely expected since the relation holds for average values of the X-ray luminosity, we notice that the stacked values of the protocluster galaxies without X-ray detection, appear to be lower than the relation measured in the field. This comparison is preliminary, since only star forming galaxies should be included. However, this is a hint that, at low mass, the accretion onto the SMBH may have been suppressed among the protocluster members, while it is enhanced at high masses. In other words, the average L_X-M_* relation may be significantly steeper within the protocluster. This issue will be explored in Pannella et al. (in preparation).



Fig. D.1. Left panel: the red points show the distribution of 12 protocluster members (including the Spiderweb Galaxy but excluding the two Compton-thick candidates) in the soft-band, intrinsic luminosity and stellar mass plane. Blue points are obtained stacking the protocluster members without X-ray detection in bins of stellar mass. Note that stacked L_X values should be considered upper limits, since, despite the formal positive detection, all the stacked values are consistent with noise within 1 or 1.5 σ. Right panel: same as the left panel but in the rest-frame hard band.